

rec.1 Primes

cmp:rec:pri:
sec

Bounded quantification and bounded minimization provide us with a good deal of machinery to show that natural functions and relations are primitive recursive. For example, consider the relation “ x divides y ”, written $x \mid y$. The relation $x \mid y$ holds if division of y by x is possible without remainder, i.e., if y is an integer multiple of x . (If it doesn’t hold, i.e., the remainder when dividing x by y is > 0 , we write $x \nmid y$.) In other words, $x \mid y$ iff for some z , $x \cdot z = y$. Obviously, any such z , if it exists, must be $\leq y$. So, we have that $x \mid y$ iff for some $z \leq y$, $x \cdot z = y$. We can define the relation $x \mid y$ by bounded existential quantification from $=$ and multiplication by

$$x \mid y \Leftrightarrow (\exists z \leq y) (x \cdot z) = y.$$

We’ve thus shown that $x \mid y$ is primitive recursive.

A natural number x is *prime* if it is neither 0 nor 1 and is only divisible by 1 and itself. In other words, prime numbers are such that, whenever $y \mid x$, either $y = 1$ or $y = x$. To test if x is prime, we only have to check if $y \mid x$ for all $y \leq x$, since if $y > x$, then automatically $y \nmid x$. So, the relation $\text{Prime}(x)$, which holds iff x is prime, can be defined by

$$\text{Prime}(x) \Leftrightarrow x \geq 2 \wedge (\forall y \leq x) (y \mid x \rightarrow y = 1 \vee y = x)$$

and is thus primitive recursive.

The primes are 2, 3, 5, 7, 11, etc. Consider the function $p(x)$ which returns the x th prime in that sequence, i.e., $p(0) = 2$, $p(1) = 3$, $p(2) = 5$, etc. (For convenience we will often write $p(x)$ as p_x ($p_0 = 2$, $p_1 = 3$, etc.)

If we had a function $\text{nextPrime}(x)$, which returns the first prime number larger than x , p can be easily defined using primitive recursion:

$$\begin{aligned} p(0) &= 2 \\ p(x+1) &= \text{nextPrime}(p(x)) \end{aligned}$$

Since $\text{nextPrime}(x)$ is the least y such that $y > x$ and y is prime, it can be easily computed by unbounded search. But it can also be defined by bounded minimization, thanks to a result due to Euclid: there is always a prime number between x and $x! + 1$.

$$\text{nextPrime}(x) = (\min y \leq x! + 1) (y > x \wedge \text{Prime}(y)).$$

This shows, that $\text{nextPrime}(x)$ and hence $p(x)$ are (not just computable but) primitive recursive.

(If you’re curious, here’s a quick proof of Euclid’s theorem. Suppose p_n is the largest prime $\leq x$ and consider the product $p = p_0 \cdot p_1 \cdot \dots \cdot p_n$ of all primes $\leq x$. Either $p + 1$ is prime or there is a prime between x and $p + 1$. Why? Suppose $p + 1$ is not prime. Then some prime number $q \mid p + 1$ where $q < p + 1$. None of the primes $\leq x$ divide $p + 1$. (By definition of p , each of the

primes $p_i \leq x$ divides p , i.e., with remainder 0. So, each of the primes $p_i \leq x$ divides $p + 1$ with remainder 1, and so $p_i \nmid p + 1$.) Hence, q is a prime $> x$ and $< p + 1$. And $p \leq x!$, so there is a prime $> x$ and $\leq x! + 1$.)

Problem rec.1. Define integer division $d(x, y)$ using bounded minimization.

Photo Credits

Bibliography