

syn.1 The De Bruijn Index

lam:syn:deb:
sec

α -Equivalence is very natural, as terms that are α -equivalent “mean the same.” In fact, it is possible to give a syntax for lambda terms which does not distinguish terms that can be α -converted to each other. The best known replaces variables by their *De Bruijn index*.

When we write $\lambda x.M$, we explicitly state that x is the parameter of the function, so that we can use x in M to refer to this parameter. In the de Bruijn index, however, parameters have no name and reference to them in the function body is denoted by a number denoting the levels of abstraction between them. For example, consider the example of $\lambda x.\lambda y.yx$: the outer abstraction is on binds the variable x ; the inner abstraction binds the variable is y ; the sub-term yx lies in the scope of the inner abstraction: there is no abstraction between y and its abstract λy , but one abstract between x and its abstract λx . Thus we write 0 1 for yx , and $\lambda.\lambda.01$ for the entire term.

Definition syn.1. De Bruijn terms are inductively defines as follows:

1. n , where n is any natural number.
2. PQ , where P and Q are both De Bruijn terms.
3. $\lambda.N$, where N is a De Bruijn term.

A formalized translation from ordinary lambda terms to De Bruijn indexed terms is as follows:

Definition syn.2.

$$\begin{aligned}F_{\Gamma}(x) &= \Gamma(x) \\F_{\Gamma}(PQ) &= F_{\Gamma}(P)F_{\Gamma}(Q) \\F_{\Gamma}(\lambda x.N) &= \lambda.F_{x,\Gamma}(N)\end{aligned}$$

where Γ is a list of variables indexed from zero, and $\Gamma(x)$ denotes the position of the variable x in Γ . For example, if Γ is x, y, z , then $\Gamma(x)$ is 0 and $\Gamma(z)$ is 2.

x, Γ denotes the list resulted from pushing x to the head of Γ ; for instance, continuing the Γ in last example, w, Γ is w, x, y, z .

Recovering a standard lambda term from a de Bruijn term is done as follows:

Definition syn.3.

$$\begin{aligned}G_{\Gamma}(n) &= \Gamma[n] \\G_{\Gamma}(PQ) &= G_{\Gamma}(P)G_{\Gamma}(Q) \\G_{\Gamma}(\lambda.N) &= \lambda x.G_{x,\Gamma}(N)\end{aligned}$$

where Γ is again a list of variables indexed from zero, and $\Gamma[n]$ denotes the variable in position n . For example, if Γ is x, y, z , then $\Gamma[1]$ is y .

The variable x in last equation is chosen to be any variable that not in Γ .

Here we give some results without proving them:

Proposition syn.4. *If $M \xrightarrow{\alpha} M'$, and Γ is any list containing $\text{FV}(M)$, then $F_{\Gamma}(M) \equiv F_{\Gamma}(M')$.*

Photo Credits

Bibliography