

choice.1 Countable Choice

It is easy to prove, without any use of Choice/Well-Ordering, that:

Lemma choice.1 (in \mathbf{Z}^-). *Every finite set has a choice function.*

Proof. Let $a = \{b_1, \dots, b_n\}$. Suppose for simplicity that each $b_i \neq \emptyset$. So there are objects c_1, \dots, c_n such that $c_1 \in b_1, \dots, c_n \in b_n$. Now by ??, the set $\{\langle b_1, c_1 \rangle, \dots, \langle b_n, c_n \rangle\}$ exists; and this is a choice function for a . \square

But matters get murkier as soon as we consider infinite sets. For example, consider this “minimal” extension to the above:

Countable Choice. Every countable set has a choice function.

This is a special case of Choice. And it transpires that this principle was invoked fairly frequently, without an obvious awareness of its use. Here are two nice examples.¹

Example 1. Here is a natural thought: for any set A , either $\omega \preceq A$, or $A \approx n$ for some $n \in \omega$. This is one way to state the intuitive idea, that every set is either finite or infinite. Cantor, and many other mathematicians, made this claim without proving it. Cautious as we are, we proved this in ??. But in that proof we were working in **ZFC**, since we were assuming that any set A can be well-ordered, and hence that $|A|$ is guaranteed to exist. That is: we explicitly assumed Choice.

In fact, [Dedekind \(1888\)](#) offered his own proof of this claim, as follows:

Theorem choice.2 (in $\mathbf{Z}^- + \text{Countable Choice}$). *For any A , either $\omega \preceq A$ or $A \approx n$ for some $n \in \omega$.*

Proof. Suppose $A \not\approx n$ for all $n \in \omega$. Then in particular for each $n < \omega$ there is subset $A_n \subseteq A$ with exactly 2^n elements. Using this sequence A_0, A_1, A_2, \dots , we define for each n :

$$B_n = A_n \setminus (A_0 \cup A_1 \cup \dots \cup A_{n-1}).$$

Now note the following

$$\begin{aligned} |A_0 \cup A_1 \cup \dots \cup A_{n-1}| &\leq |A_0| + |A_1| + \dots + |A_{n-1}| \\ &= 1 + 2 + \dots + 2^{n-1} \\ &= 2^n - 1 \\ &< 2^n = |A_n| \end{aligned}$$

Hence each B_n has at least one member, c_n . Moreover, the B_n s are pairwise disjoint; so if $c_n = c_m$ then $n = m$. But every $c_n \in A$. So the function $f(n) = c_n$ is an injection $\omega \rightarrow A$. \square

¹Due to [Potter \(2004, §9.4\)](#) and Luca Incurvati.

Dedekind did not flag that he had used Countable Choice. But, did *you* spot its use? Look again. (Really: Look again.)

The proof used Countable Choice twice. We used it once, to obtain our sequence of sets A_0, A_1, A_2, \dots . We then used it again to select our elements c_n from each B_n . Moreover, this use of Choice is ineliminable. Cohen (1966, p. 138) proved that the result fails if we have no version of Choice. That is: it is consistent with **ZF** that there are sets which are *incomparable* with ω .

Example 2. In 1878, Cantor stated that a countable union of countable sets is countable. He did not present a proof, perhaps indicating that he took the proof to be obvious. Now, cautious as we are, we proved a more general version of this result in ???. But our proof explicitly assumed Choice. And even the proof of the less general result requires Countable Choice.

Theorem choice.3 (in $\mathbf{Z}^- + \text{Countable Choice}$). *If A_n is countable for each $n \in \omega$, then $\bigcup_{n < \omega} A_n$ is countable.*

Proof. Without loss of generality, suppose that each $A_n \neq \emptyset$. So for each $n \in \omega$ there is a surjection $f_n: \omega \rightarrow A_n$. Define $f: \omega \times \omega \rightarrow \bigcup_{n < \omega} A_n$ by $f(m, n) = f_n(m)$. The result follows because $\omega \times \omega$ is countable (??) and f is a surjection. \square

Did you spot the use of the Countable Choice? It is used to choose our sequence of functions f_0, f_1, f_2, \dots .² And again, the result fails in the absence of any Choice principle. Specifically, Feferman and Levy (1963) proved that it is consistent with **ZF** that a countable union of countable sets has cardinality \beth_1 . But here is a much funnier statement of the point, from Russell:

This is illustrated by the millionaire who bought a pair of socks whenever he bought a pair of boots, and never at any other time, and who had such a passion for buying both that at last he had \aleph_0 pairs of boots and \aleph_0 pairs of socks... Among boots we can distinguish right and left, and therefore we can make a selection of one out of each pair, namely, we can choose all the right boots or all the left boots; but with socks no such principle of selection suggests itself, and we cannot be sure, unless we assume the multiplicative axiom [i.e., in effect Choice], that there is any class consisting of one sock out of each pair. (Russell, 1919, p. 126)

In short, some form of Choice is needed to prove the following: If you have countably many pairs of socks, then you have (only) countably many socks. And in fact, without Countable Choice (or something equivalent), a countable union of countable sets can fail to be countable.

The moral is that Countable Choice was used repeatedly, without much awareness of its users. The philosophical question is: How could we *justify* Countable Choice?

²A similar use of Choice occurred in ??? when we “fixed a bijection f_β ” for each $\beta \in \alpha$.

An attempt at an intuitive justification might invoke an appeal to a super-task. Suppose we make the first choice in $1/2$ a minute, our second choice in $1/4$ a minute, \dots , our n -th choice in $1/2^n$ a minute, \dots . Then within 1 minute, we will have made an ω -sequence of choices, and defined a choice function.

But what, really, could such a thought-experiment tell us? For a start, it relies upon taking this idea of “choosing” rather literally. For another, it seems to bind up mathematics in metaphysical possibility.

More important: it is not going to give us any justification for Choice *tout court*, rather than *mere* Countable Choice. For if we need *every* set to have a choice function, then we will need to be able to perform a “supertask of arbitrary ordinal length”. Bluntly, that idea is laughable.

Photo Credits

Bibliography

Cantor, Georg. 1878. Ein Beitrag zur Mannigfaltigkeitslehre. *Journal für die reine und angewandte Mathematik* 84: 242–58.

Cohen, Paul J. 1966. *Set Theory and the Continuum Hypothesis*. Reading, MA: Benjamin.

Dedekind, Richard. 1888. *Was sind und was sollen die Zahlen?* Braunschweig: Vieweg.

Feferman, Solomon and Azriel Levy. 1963. Independence results in set theory by Cohen’s method II. *Notices of the American Mathematical Society* 10: 593.

Potter, Michael. 2004. *Set Theory and its Philosophy*. Oxford: Oxford University Press.

Russell, Bertrand. 1919. *Introduction to Mathematical Philosophy*. London: Allen & Unwin.