rel.1 Orders

sfr:rel:ord:sec Many of our comparisons involve describing some objects as being "less than", "equal to", or "greater than" other objects, in a certain respect. These involve *order* relations. But there are different kinds of order relations. For instance, some require that any two objects be comparable, others don't. Some include identity (like $\leq$) and some exclude it (like $<$). It will help us to have a taxonomy here. explanation

Definition rel.1 (Preorder). A relation which is both reflexive and transitive is called a *preorder.*

Definition rel.2 (Partial order). A preorder which is also anti-symmetric is called a *partial order.*

Definition rel.3 (Linear order). A partial order which is also connected is called a *total order* or *linear order.*

Every linear order is also a partial order, and every partial order is also a preorder, but the converses don't hold.

Example rel.4. Every linear order is also a partial order, and every partial order is also a preorder, but the converses don't hold. The universal relation on A is a preorder, since it is reflexive and transitive. But, if A has more than one element, the universal relation is not anti-symmetric, and so not a partial order.

Example rel.5. Consider the *no longer than* relation $\preccurlyeq$ on $\mathbb{B}^*$: $x \preccurlyeq y$ iff $\text{len}(x) \leq \text{len}(y)$. This is a preorder (reflexive and transitive), and even connected, but not a partial order, since it is not anti-symmetric. For instance, $01 \preccurlyeq 10$ and $10 \preccurlyeq 01$, but $01 \neq 10$.

Example rel.6. An important partial order is the relation $\subseteq$ on a set of sets. This is not in general a linear order, since if $a \neq b$ and we consider $\wp(\{a, b\}) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$, we see that $\{a\} \nsubseteq \{b\}$ and $\{a\} \neq \{b\}$ and $\{b\} \nsubseteq \{a\}$.

Example rel.7. The relation of *divisibility without remainder* gives us a partial order which isn't a linear order. For integers n, m, we write $n \mid m$ to mean n (evenly) divides m, i.e., iff there is some integer k so that $m = kn$. On $\mathbb{N}$, this is a partial order, but not a linear order: for instance, $2 \nmid 3$ and also $3 \nmid 2$. Considered as a relation on $\mathbb{Z}$, divisibility is only a preorder since it is not anti-symmetric: $1 \mid -1$ and $-1 \mid 1$ but $1 \neq -1$.

Definition rel.8 (Strict order). A *strict order* is a relation which is irreflexive, asymmetric, and transitive.

Definition rel.9 (Strict linear order). A strict order which is also connected is called a *strict linear order.*

Example rel.10. $\leq$ is the linear order corresponding to the strict linear order $<$. $\subseteq$ is the partial order corresponding to the strict order $\subsetneq$.

Definition rel.11 (Total order). A strict order which is also connected is called a *total order*. This is also sometimes called a *strict linear order*. sfr:rel:ord: def:strictlinearorder

Any strict order R on A can be turned into a partial order by adding the diagonal Id_A, i.e., adding all the pairs $\langle x, x \rangle$. (This is called the *reflexive closure* of R.) Conversely, starting from a partial order, one can get a strict order by removing Id_A. These next two results make this precise.

Proposition rel.12. *If R is a strict order on A, then $R^+ = R \cup \mathrm{Id}_A$ is a partial order. Moreover, if R is total, then R^+ is a linear order.* *sfr:rel:ord: prop:stricttopartial*

Proof. Suppose R is a strict order, i.e., $R \subseteq A^2$ and R is irreflexive, asymmetric, and transitive. Let $R^+ = R \cup \mathrm{Id}_A$. We have to show that R^+ is reflexive, antisymmetric, and transitive.

R^+ is clearly reflexive, since $\langle x, x \rangle \in \mathrm{Id}_A \subseteq R^+$ for all $x \in A$.

To show R^+ is antisymmetric, suppose for reductio that R^+xy and R^+yx but $x \neq y$. Since $\langle x, y \rangle \in R \cup \mathrm{Id}_X$, but $\langle x, y \rangle \notin \mathrm{Id}_X$, we must have $\langle x, y \rangle \in R$, i.e., Rxy. Similarly, Ryx. But this contradicts the assumption that R is asymmetric.

To establish transitivity, suppose that R^+xy and R^+yz. If both $\langle x, y \rangle \in R$ and $\langle y, z \rangle \in R$, then $\langle x, z \rangle \in R$ since R is transitive. Otherwise, either $\langle x, y \rangle \in \mathrm{Id}_X$, i.e., $x = y$, or $\langle y, z \rangle \in \mathrm{Id}_X$, i.e., $y = z$. In the first case, we have that R^+yz by assumption, $x = y$, hence R^+xz. Similarly in the second case. In either case, R^+xz, thus, R^+ is also transitive.

Concerning the "moreover" clause, suppose R is a total order, i.e., that R is connected. So for all $x \neq y$, either Rxy or Ryx, i.e., either $\langle x, y \rangle \in R$ or $\langle y, x \rangle \in R$. Since $R \subseteq R^+$, this remains true of R^+, so R^+ is connected as well. $\square$

Proposition rel.13. *If R is a partial order on X, then $R^- = R \setminus \mathrm{Id}_X$ is a strict order. Moreover, if R is linear, then R^- is total.* *sfr:rel:ord: prop:partialtostrict*

Proof. This is left as an exercise. $\square$

Problem rel.1. Give a proof of Proposition rel.13.

Example rel.14. $\leq$ is the linear order corresponding to the total order $<$. $\subseteq$ is the partial order corresponding to the strict order $\subsetneq$.

The following simple result which establishes that total orders satisfy an extensionality-like property:

Proposition rel.15. *If $<$ totally orders A, then:* *sfr:rel:ord: prop:extensionality-totalorders*

$$(\forall a, b \in A)((\forall x \in A)(x < a \leftrightarrow x < b) \rightarrow a = b)$$

Proof. Suppose $(\forall x \in A)(x < a \leftrightarrow x < b)$. If $a < b$, then $a < a$, contradicting the fact that $<$ is irreflexive; so $a \not< b$. Exactly similarly, $b \not< a$. So $a = b$, as $<$ is connected. $\square$

Photo Credits

Bibliography