

Figure 1: The union $X \cup Y$ of two sets is set of elements of X together with those of Y.

set.1 Unions and Intersections

sfr:set:uni:sec We can define new sets by abstraction, and the property used to define the new set can mention sets we've already defined. So for instance, if X and Y are sets, the set $\{x : x \in X \lor x \in Y\}$ defines a set which consists of all those objects which are elements of either X or Y, i.e., it's the set that combines the elements of X and Y. This operation on sets—combining them—is very useful and common, and so we give it a name and a symbol. explanation

Definition set.1 (Union)**.** The *union* of two sets X and Y, written $X \cup Y$, is the set of all things which are elements of X, Y, or both.

$$X \cup Y = \{x : x \in X \lor x \in Y\}$$

Example set.2. Since the multiplicity of elements doesn't matter, the union of two sets which have an element in common contains that element only once, e.g., $\{a, b, c\} \cup \{a, 0, 1\} = \{a, b, c, 0, 1\}$.

The union of a set and one of its subsets is just the bigger set: $\{a, b, c\} \cup \{a\} = \{a, b, c\}$.

The union of a set with the empty set is identical to the set: $\{a, b, c\} \cup \emptyset = \{a, b, c\}$.

Problem set.1. Prove rigorously that if $X \subseteq Y$, then $X \cup Y = Y$.

The operation that forms the set of all elements that X and Y have in common is called their *intersection*. explanation

Definition set.3 (Intersection)**.** The *intersection* of two sets X and Y, written $X \cap Y$, is the set of all things which are elements of both X and Y.

$$X \cap Y = \{x : x \in X \land x \in Y\}$$



Figure 2: The intersection $X \cap Y$ of two sets is the set of elements they have in common.

Two sets are called *disjoint* if their intersection is empty. This means they have no elements in common.

Example set.4. If two sets have no elements in common, their intersection is empty: $\{a, b, c\} \cap \{0, 1\} = \emptyset$.

If two sets do have elements in common, their intersection is the set of all those: $\{a, b, c\} \cap \{a, b, d\} = \{a, b\}$.

The intersection of a set with one of its subsets is just the smaller set: $\{a, b, c\} \cap \{a, b\} = \{a, b\}$.

The intersection of any set with the empty set is empty: $\{a, b, c\} \cap \emptyset = \emptyset$.

Problem set.2. Prove rigorously that if $X \subseteq Y$, then $X \cap Y = X$.

explanation

We can also form the union or intersection of more than two sets. An elegant way of dealing with this in general is the following: suppose you collect all the sets you want to form the union (or intersection) of into a single set. Then we can define the union of all our original sets as the set of all objects which belong to at least one element of the set, and the intersection as the set of all objects which belong to every element of the set.

Definition set.5. If Z is a set of sets, then $\bigcup Z$ is the set of elements of elements of Z:

$$\bigcup Z = \{x : x \text{ belongs to an element of } Z\}, \text{ i.e.,}$$
$$\bigcup Z = \{x : \text{there is a } Y \in Z \text{ so that } x \in Y\}$$

Definition set.6. If Z is a set of sets, then $\bigcap Z$ is the set of objects which all elements of Z have in common:

$$\bigcap Z = \{x : x \text{ belongs to every element of } Z\}, \text{ i.e.,}$$
$$\bigcap Z = \{x : \text{for all } Y \in Z, x \in Y\}$$



Figure 3: The difference $X \setminus Y$ of two sets is the set of those elements of X which are not also elements of Y.

Example set.7. Suppose $Z = \{\{a, b\}, \{a, d, e\}, \{a, d\}\}$. Then $\bigcup Z = \{a, b, d, e\}$ and $\bigcap Z = \{a\}$.

We could also do the same for a sequence of sets X_1, X_2, ...

$$\bigcup_i X_i = \{x : x \text{ belongs to one of the } X_i\}$$

$$\bigcap_i X_i = \{x : x \text{ belongs to every } X_i\}.$$

Definition set.8 (Difference)**.** The *difference* $X \setminus Y$ is the set of all elements of X which are not also elements of Y, i.e.,

$$X \setminus Y = \{x : x \in X \text{ and } x \notin Y\}.$$

Photo Credits

Bibliography