

## int.1 First-Order Logic

fol:int:fol:  
sec

You are probably familiar with first-order logic from your first introduction to formal logic.<sup>1</sup> You may know it as “quantificational logic” or “predicate logic.” First-order logic, first of all, is a formal language. That means, it has a certain vocabulary, and its expressions are strings from this vocabulary. But not every string is permitted. There are different kinds of permitted expressions: terms, formulas, and sentences. We are mainly interested in sentences of first-order logic: they provide us with a formal analogue of sentences of English, and about them we can ask the questions a logician typically is interested in. For instance:

- Does  $\psi$  follow from  $\varphi$  logically?
- Is  $\varphi$  logically true, logically false, or contingent?
- Are  $\varphi$  and  $\psi$  equivalent?

These questions are primarily questions about the “meaning” of sentences of first-order logic. For instance, a philosopher would analyze the question of whether  $\psi$  follows logically from  $\varphi$  as asking: is there a case where  $\varphi$  is true but  $\psi$  is false ( $\psi$  doesn’t follow from  $\varphi$ ), or does every case that makes  $\varphi$  true also make  $\psi$  true ( $\psi$  does follow from  $\varphi$ )? But we haven’t been told yet what a “case” is—that is the job of *semantics*. The semantics of first-order logic provides a mathematically precise model of the philosopher’s intuitive idea of “case,” and also—and this is important—of what it is for a sentence  $\varphi$  to be true in a case. We call the mathematically precise model that we will develop a structure. The relation which makes “true in” precise, is called the relation of satisfaction. So what we will define is “ $\varphi$  is satisfied in  $\mathfrak{M}$ ” (in symbols:  $\mathfrak{M} \models \varphi$ ) for sentences  $\varphi$  and structures  $\mathfrak{M}$ . Once this is done, we can also give precise definitions of the other semantical terms such as “follows from” or “is logically true.” These definitions will make it possible to settle, again with mathematical precision, whether, e.g.,  $\forall x (\varphi(x) \rightarrow \psi(x))$ ,  $\exists x \varphi(x) \models \exists x \psi(x)$ . The answer will, of course, be “yes.” If you’ve already been trained to symbolize sentences of English in first-order logic, you will recognize this as, e.g., the symbolizations of, say, “All ants are insects, there are ants, therefore there are insects.” That is obviously a valid argument, and so our mathematical model of “follows from” for our formal language should give the same answer.

Another topic you probably remember from your first introduction to formal logic is that there are *derivations*. If you have taken a first formal logic course, your instructor will have made you practice finding such derivations, perhaps even a derivation that shows that the above entailment holds. There are many different ways to give derivations: you may have done something called “natural deduction” or “truth trees,” but there are many others. The purpose of derivation systems is to provide tools using which the logicians’

---

<sup>1</sup>In fact, we more or less assume you are! If you’re not, you could review a more elementary textbook, such as *forall x* (Magnus et al., 2021).

questions above can be answered: e.g., a natural deduction **derivation** in which  $\forall x (\varphi(x) \rightarrow \psi(x))$  and  $\exists x \varphi(x)$  are premises and  $\exists x \psi(x)$  is the conclusion (last line) *verifies* that  $\exists x \psi(x)$  logically follows from  $\forall x (\varphi(x) \rightarrow \psi(x))$  and  $\exists x \varphi(x)$ .

But why is that? On the face of it, **derivation** systems have nothing to do with semantics: giving a formal **derivation** merely involves arranging symbols in certain rule-governed ways; they don't mention "cases" or "true in" at all. The connection between **derivation** systems and semantics has to be established by a meta-logical investigation. What's needed is a mathematical proof, e.g., that a formal **derivation** of  $\exists x \psi(x)$  from premises  $\forall x (\varphi(x) \rightarrow \psi(x))$  and  $\exists x \varphi(x)$  is possible, if, and only if,  $\forall x (\varphi(x) \rightarrow \psi(x))$  and  $\exists x \varphi(x)$  together entails  $\exists x \psi(x)$ . Before this can be done, however, a lot of painstaking work has to be carried out to get the definitions of syntax and semantics correct.

## Photo Credits

## Bibliography

Magnus, P. D., Tim Button, J. Robert Loftis, Aaron Thomas-Bolduc, Robert Trueman, and Richard Zach. 2021. *forall x: Calgary. An Introduction to Formal Logic*. Calgary: Open Logic Project, f21 ed. URL <https://forallx.openlogicproject.org/>.