

## int.1 Satisfaction

fol:int:sat:  
sec

We can already skip ahead to the semantics of first-order logic once we know what **formulas** are: here, the basic definition is that of a **structure**. For our simple language, a **structure**  $\mathfrak{M}$  has just three components: a non-empty set  $|\mathfrak{M}|$  called the *domain*, what  $a$  picks out in  $\mathfrak{M}$ , and what  $P$  is true of in  $\mathfrak{M}$ . The object picked out by  $a$  is denoted  $a^{\mathfrak{M}}$  and the set of things  $P$  is true of by  $P^{\mathfrak{M}}$ . A **structure**  $\mathfrak{M}$  consists of just these three things:  $|\mathfrak{M}|$ ,  $a^{\mathfrak{M}} \in |\mathfrak{M}|$  and  $P^{\mathfrak{M}} \subseteq |\mathfrak{M}|$ . The general case will be more complicated, since there will be many **predicate symbols** and **constant symbols**, the **constant symbols** can have more than one place, and there will also be **function symbols**.

This is enough to give a definition of satisfaction for **formulas** that don't contain **variables**. The idea is to give an inductive definition that mirrors the way we have defined **formulas**. We specify when an atomic formula is satisfied in  $\mathfrak{M}$ , and then when, e.g.,  $\neg\varphi$  is satisfied in  $\mathfrak{M}$  on the basis of whether or not  $\varphi$  is satisfied in  $\mathfrak{M}$ . E.g., we could define:

1.  $P(a)$  is satisfied in  $\mathfrak{M}$  iff  $a^{\mathfrak{M}} \in P^{\mathfrak{M}}$ .
2.  $\neg\varphi$  is satisfied in  $\mathfrak{M}$  iff  $\varphi$  is not satisfied in  $\mathfrak{M}$ .
3.  $(\varphi \wedge \psi)$  is satisfied in  $\mathfrak{M}$  iff  $\varphi$  is satisfied in  $\mathfrak{M}$ , and  $\psi$  is satisfied in  $\mathfrak{M}$  as well.

Let's say that  $|\mathfrak{M}| = \{0, 1, 2\}$ ,  $a^{\mathfrak{M}} = 1$ , and  $P^{\mathfrak{M}} = \{1, 2\}$ . This definition would tell us that  $P(a)$  is satisfied in  $\mathfrak{M}$  (since  $a^{\mathfrak{M}} = 1 \in \{1, 2\} = P^{\mathfrak{M}}$ ). It tells us further that  $\neg P(a)$  is not satisfied in  $\mathfrak{M}$ , and that in turn  $\neg\neg P(a)$  is and  $(\neg P(a) \wedge P(a))$  is not satisfied, and so on.

The trouble comes when we want to give a definition for the quantifiers: we'd like to say something like, " $\exists v_0 P(v_0)$  is satisfied iff  $P(v_0)$  is satisfied." But the **structure**  $\mathfrak{M}$  doesn't tell us what to do about **variables**. What we actually want to say is that  $P(v_0)$  is satisfied *for some value of*  $v_0$ . To make this precise we need a way to assign **elements** of  $|\mathfrak{M}|$  not just to  $a$  but also to  $v_0$ . To this end, we introduce **variable assignments**. A **variable assignment** is simply a function  $s$  that maps **variables** to **elements** of  $|\mathfrak{M}|$  (in our example, to one of 1, 2, or 3). Since we don't know beforehand which **variables** might appear in a **formula** we can't limit which **variables**  $s$  assigns values to. The simple solution is to require that  $s$  assigns values to *all* **variables**  $v_0, v_1, \dots$ . We'll just use only the ones we need.

Instead of defining satisfaction of **formulas** just relative to a **structure**, we'll define it relative to a **structure**  $\mathfrak{M}$  and a **variable assignment**  $s$ , and write  $\mathfrak{M}, s \models \varphi$  for short. Our definition will now include an additional clause to deal with atomic **formulas** containing **variables**:

1.  $\mathfrak{M}, s \models P(a)$  iff  $a^{\mathfrak{M}} \in P^{\mathfrak{M}}$ .
2.  $\mathfrak{M}, s \models P(v_i)$  iff  $s(v_i) \in P^{\mathfrak{M}}$ .
3.  $\mathfrak{M}, s \models \neg\varphi$  iff not  $\mathfrak{M}, s \models \varphi$ .

4.  $\mathfrak{M}, s \models (\varphi \wedge \psi)$  iff  $\mathfrak{M}, s \models \varphi$  and  $\mathfrak{M}, s \models \psi$ .

Ok, this solves one problem: we can now say when  $\mathfrak{M}$  satisfies  $P(v_0)$  for the value  $s(v_0)$ . To get the definition right for  $\exists v_0 P(v_0)$  we have to do one more thing: We want to have that  $\mathfrak{M}, s \models \exists v_0 P(v_0)$  iff  $\mathfrak{M}, s' \models P(v_0)$  for *some* way  $s'$  of assigning a value to  $v_0$ . But the value assigned to  $v_0$  does not necessarily have to be the value that  $s(v_0)$  picks out. We'll introduce a notation for that: if  $m \in |\mathfrak{M}|$ , then we let  $s[m/v_0]$  be the assignment that is just like  $s$  (for all **variables** other than  $v_0$ ), except to  $v_0$  it assigns  $m$ . Now our definition can be:

5.  $\mathfrak{M}, s \models \exists v_i \varphi$  iff  $\mathfrak{M}, s[m/v_i] \models \varphi$  for some  $m \in |\mathfrak{M}|$ .

Does it work out? Let's say we let  $s(v_i) = 0$  for all  $i \in \mathbb{N}$ .  $\mathfrak{M}, s \models \exists v_0 P(v_0)$  iff there is an  $m \in |\mathfrak{M}|$  so that  $\mathfrak{M}, s[m/v_0] \models P(v_0)$ . And there is: we can choose  $m = 1$  or  $m = 2$ . Note that this is true even if the value  $s(v_0)$  assigned to  $v_0$  by  $s$  itself—in this case, 0—doesn't do the job. We have  $\mathfrak{M}, s[1/v_0] \models P(v_0)$  but not  $\mathfrak{M}, s \models P(v_0)$ .

If this looks confusing and cumbersome: it is. But the added complexity is required to give a precise, inductive definition of satisfaction for all **formulas**, and we need something like it to precisely define the semantic notions. There are other ways of doing it, but they are all equally (in)elegant.

## Photo Credits

## Bibliography