

set.1 Cantor on the Line and the Plane

his:set:cantorplane:
sec

Some of the circumstances surrounding the proof of Schröder-Bernstein tie in with the history we discussed in ???. Recall that, in 1877, Cantor proved that there are exactly as many points on a square as on one of its sides. Here, we will present his (first attempted) proof.

Let L be the unit line, i.e., the set of points $[0, 1]$. Let S be the unit square, i.e., the set of points $L \times L$. In these terms, Cantor proved that $L \approx S$. He wrote a note to Dedekind, essentially containing the following argument.

his:set:cantorplane:
thm:cantorplane

Theorem set.1. $L \approx S$

Proof: first part.. Fix $a, b \in L$. Write them in binary notation, so that we have infinite sequences of 0s and 1s, a_1, a_2, \dots , and b_1, b_2, \dots , such that:

$$\begin{aligned} a &= 0.a_1a_2a_3a_4\dots \\ b &= 0.b_1b_2b_3b_4\dots \end{aligned}$$

Now consider the function $f: S \rightarrow L$ given by

$$f(a, b) = 0.a_1b_1a_2b_2a_3b_3a_4b_4\dots$$

Now f is an **injection**, since if $f(a, b) = f(c, d)$, then $a_n = c_n$ and $b_n = d_n$ for all $n \in \mathbb{N}$, so that $a = c$ and $b = d$. \square

Unfortunately, as Dedekind pointed out to Cantor, this does not answer the original question. Consider $0.\dot{1}0 = 0.10101010\dots$. We need that $f(a, b) = 0.\dot{1}0$, where:

$$\begin{aligned} a &= 0.\dot{1}1 = 0.111111\dots \\ b &= 0 \end{aligned}$$

But $a = 0.\dot{1}1 = 1$. So, when we say “write a and b in binary notation”, we have to choose *which* notation to use; and, since f is to be a *function*, we can use only *one* of the two possible notations. But if, for example, we use the simple notation, and write a as “1.000...”, then we have no pair $\langle a, b \rangle$ such that $f(a, b) = 0.\dot{1}0$.

To summarise: Dedekind pointed out that, given the possibility of certain recurring decimal expansions, Cantor’s function f is an **injection** but *not* a **surjection**. So Cantor has shown only that $S \preceq L$ and *not* that $S \approx L$.

Cantor wrote back to Dedekind almost immediately, essentially suggesting that the proof could be completed as follows:

Proof: completed.. So, we have shown that $S \preceq L$. But there is obviously an **injection** from L to S : just lay the line flat along one side of the square. So $L \preceq S$ and $S \preceq L$. By Schröder-Bernstein (??), $L \approx S$. \square

But of course, Cantor could not complete the last line in these terms, for the Schröder-Bernstein Theorem was not yet proved. Indeed, although Cantor would subsequently formulate this as a general conjecture, it was not satisfactorily proved until 1897. (And so, later in 1877, Cantor offered a different proof of [Theorem set.1](#), which did not go via Schröder–Bernstein.)

Photo Credits

Bibliography