## met.1 Second-order Logic is not Compact

 $\begin{array}{c} \mathbf{sol} : \mathbf{met} : \mathbf{com} \colon \\ \mathbf{sec} \end{array}$ 

Call a set of sentences  $\Gamma$  finitely satisfiable if every one of its finite subsets is satisfiable. First-order logic has the property that if a set of sentences  $\Gamma$  is finitely satisfiable, it is satisfiable. This property is called *compactness*. It has an equivalent version involving entailment: if  $\Gamma \vDash \varphi$ , then already  $\Gamma_0 \vDash \varphi$  for some finite subset  $\Gamma_0 \subseteq \Gamma$ . In this version it is an immediate corollary of the completeness theorem: for if  $\Gamma \vDash \varphi$ , by completeness  $\Gamma \vdash \varphi$ . But a derivation can only make use of finitely many sentences of  $\Gamma$ .

Compactness is not true for second-order logic. There are sets of second-order sentences that are finitely satisfiable but not satisfiable, and that entail some  $\varphi$  without a finite subset entailing  $\varphi$ .

 $sol:met:com: \\ thm:sol-undecidable$ 

Theorem met.1. Second-order logic is not compact.

Proof. Recall that

Inf 
$$\equiv \exists u (\forall x \forall y (u(x) = u(y) \rightarrow x = y) \land \exists y \forall x y \neq u(x))$$

is satisfied in a structure iff its domain is infinite. Let  $\varphi^{\geq n}$  be a sentence that asserts that the domain has at least n elements, e.g.,

$$\varphi^{\geq n} \equiv \exists x_1 \dots \exists x_n (x_1 \neq x_2 \land x_1 \neq x_3 \land \dots \land x_{n-1} \neq x_n).$$

Consider the set of sentences

$$\Gamma = {\neg \operatorname{Inf}, \varphi^{\geq 1}, \varphi^{\geq 2}, \varphi^{\geq 3}, \dots}.$$

It is finitely satisfiable, since for any finite subset  $\Gamma_0 \subseteq \Gamma$  there is some k so that  $\varphi^{\geq k} \in \Gamma$  but no  $\varphi^{\geq n} \in \Gamma$  for n > k. If  $|\mathfrak{M}|$  has k elements,  $\mathfrak{M} \models \Gamma_0$ . But,  $\Gamma$  is not satisfiable: if  $\mathfrak{M} \models \neg \text{Inf}$ ,  $|\mathfrak{M}|$  must be finite, say, of size k. Then  $\mathfrak{M} \nvDash \varphi^{\geq k+1}$ .

**Problem met.1.** Give an example of a set  $\Gamma$  and a sentence  $\varphi$  so that  $\Gamma \vDash \varphi$  but for every finite subset  $\Gamma_0 \subseteq \Gamma$ ,  $\Gamma_0 \nvDash \varphi$ .

**Photo Credits** 

Bibliography