

## ord-arithmetic.1 Using Ordinal Addition

Using addition on the ordinals, we can explicitly calculate the ranks of various sets, in the sense of ??:

**Lemma ord-arithmetic.1.** *If  $\text{rank}(A) = \alpha$  and  $\text{rank}(B) = \beta$ , then:*

1.  $\text{rank}(\wp(A)) = \alpha + 1$
2.  $\text{rank}(\{A, B\}) = \max(\alpha, \beta) + 1$
3.  $\text{rank}(A \cup B) = \max(\alpha, \beta)$
4.  $\text{rank}(\langle A, B \rangle) = \max(\alpha, \beta) + 2$
5.  $\text{rank}(A \times B) \leq \max(\alpha, \beta) + 2$
6.  $\text{rank}(\bigcup A) = \alpha$  when  $\alpha$  is empty or a limit;  $\text{rank}(\bigcup A) = \gamma$  when  $\alpha = \gamma + 1$

*Proof.* Throughout, we invoke ?? repeatedly.

(1). If  $x \subseteq A$  then  $\text{rank}(x) \leq \text{rank}(A)$ . So  $\text{rank}(\wp(A)) \leq \alpha + 1$ . Since  $A \in \wp(A)$  in particular,  $\text{rank}(\wp(A)) = \alpha + 1$ .

(2). By ??

(3). By ??.

(4). By (2), twice.

(5). Note that  $A \times B \subseteq \wp(\wp(A \cup B))$ , and invoke (4).

(6). If  $\alpha = \gamma + 1$ , there is some  $c \in A$  with  $\text{rank}(c) = \gamma$ , and no element of  $A$  has higher rank; so  $\text{rank}(\bigcup A) = \gamma$ . If  $\alpha$  is a limit ordinal, then  $A$  has elements with rank arbitrarily close to (but strictly less than)  $\alpha$ , so that  $\bigcup A$  also has elements with rank arbitrarily close to (but strictly less than)  $\alpha$ , so that  $\text{rank}(\bigcup A) = \alpha$ .  $\square$

We leave it as an exercise to show why (5) involves an inequality.

**Problem ord-arithmetic.1.** Produce sets  $A$  and  $B$  such that  $\text{rank}(A \times B) = \max(\text{rank}(A), \text{rank}(B))$ . Produce sets  $A$  and  $B$  such that  $\text{rank}(A \times B) > \max(\text{rank}(A), \text{rank}(B))$ .  
2. Are any other ranks possible?

We are also now in a position to show that several reasonable notions of what it might mean to describe an ordinal as “finite” or “infinite” coincide:

**Lemma ord-arithmetic.2.** *For any ordinal  $\alpha$ , the following are equivalent:*

1.  $\alpha \notin \omega$ , i.e.,  $\alpha$  is not a natural number
2.  $\omega \leq \alpha$
3.  $1 + \alpha = \alpha$
4.  $\alpha \approx \alpha + 1$ , i.e.,  $\alpha$  and  $\alpha + 1$  are equinumerous

5.  $\alpha$  is Dedekind infinite

[sth:ord-arithmetic:using-addition](#)  
[ord:infinite](#)

So we have five provably equivalent ways to understand what it takes for an ordinal to be (in)finite.

*Proof.* (1)  $\Rightarrow$  (2). By Trichotomy.

(2)  $\Rightarrow$  (3). Fix  $\alpha \geq \omega$ . By Transfinite Induction, there is some least ordinal  $\gamma$  (possibly 0) such that there is a limit ordinal  $\beta$  with  $\alpha = \beta + \gamma$ . Now:

$$1 + \alpha = 1 + (\beta + \gamma) = (1 + \beta) + \gamma = \operatorname{lsub}_{\delta < \beta}(1 + \delta) + \gamma = \beta + \gamma = \alpha.$$

(3)  $\Rightarrow$  (4). There is clearly a bijection  $f: (\alpha \sqcup 1) \rightarrow (1 \sqcup \alpha)$ . If  $1 + \alpha = \alpha$ , there is an isomorphism  $g: (1 \sqcup \alpha) \rightarrow \alpha$ . Now consider  $g \circ f$ .

(4)  $\Rightarrow$  (5). If  $\alpha \approx \alpha + 1$ , there is a bijection  $f: (\alpha \sqcup 1) \rightarrow \alpha$ . Define  $g(\gamma) = f(\gamma, 0)$  for each  $\gamma < \alpha$ ; this injection witnesses that  $\alpha$  is Dedekind infinite, since  $f(0, 1) \in \alpha \setminus \operatorname{ran}(g)$ .

(5)  $\Rightarrow$  (1). This is ??.

□

## Photo Credits

## Bibliography