

z.1 Appendix: Closure, Comprehension, and Intersection

sth:z:arbintersections: sec In ??, we suggested that you should look back through the naïve work of ?? and check that it can be carried out in \mathbf{Z}^- . If you followed that advice, *one* point might have tripped you up: the use of *intersection* in Dedekind’s treatment of *closures*.

Recall from ?? that

$$\text{clo}_f(o) = \bigcap \{X : o \in X \text{ and } X \text{ is } f\text{-closed}\}.$$

The general shape of this is a definition of the form:

$$C = \bigcap \{X : \varphi(X)\}.$$

But this should ring alarm bells: since Naïve Comprehension fails, there is no guarantee that $\{X : \varphi(X)\}$ exists. It looks dangerously, then, like such definitions are *cheating*.

Fortunately, they are not cheating; or rather, if they *are* cheating as they stand, then we can engage in some honest toil to render them kosher. That honest toil was foreshadowed in ??, when we explained why $\bigcap A$ exists for any $A \neq \emptyset$. But we will spell it out explicitly.

Given Extensionality, if we attempt to define C as $\bigcap \{X : \varphi(X)\}$, all we are really asking is for an object C which obeys the following:

$$\forall x(x \in C \leftrightarrow \forall X(\varphi(X) \rightarrow x \in X)) \tag{*}$$

sth:z:arbintersections: bicondelimarbintersection

Now, suppose there is *some* set, S , such that $\varphi(S)$. Then to deliver eq. (*), we can simply define C using *Separation*, as follows:

$$C = \{x \in S : \forall X(\varphi(X) \rightarrow x \in X)\}.$$

We leave it as an exercise to check that this definition yields eq. (*), as desired. And this general strategy will allow us to circumvent any apparent use of Naïve Comprehension in defining intersections. In the particular case which got us started on this line of thought, namely that of $\text{clo}_f(o)$, here is how that would work. We began the proof of ?? by noting that $o \in \text{ran}(f) \cup \{o\}$ and that $\text{ran}(f) \cup \{o\}$ is f -closed. So, we can define what we want thus:

$$\text{clo}_f(o) = \{x \in \text{ran}(f) \cup \{o\} : (\forall X \ni o)(X \text{ is } f\text{-closed} \rightarrow x \in X)\}.$$

Photo Credits

Bibliography