

-NoValue-

thy.1 Introduction

cmp:thy:int:
sec

The branch of logic known as *computability theory* deals with issues having to do with the computability, or relative computability, of functions and sets. It is evidence of Kleene's influence that the subject used to be known as *recursion theory*, and today, both names are commonly used.

Let us call a function $f: \mathbb{N} \rightarrow \mathbb{N}$ *partial computable* if it can be computed in some model of computation. If f is total we will simply say that f is *computable*. A relation R with computable characteristic function χ_R is also called computable. If f and g are partial functions, we will write $f(x) \downarrow$ to mean that f is defined at x , O.s., x is in the domain of f ; and $f(x) \uparrow$ to mean the opposite, O.s., that f is not defined at x . We will use $f(x) \simeq g(x)$ to mean that either $f(x)$ and $g(x)$ are both undefined, or they are both defined and equal.

One can explore the theory of computability without having to refer to a specific model of computation. To do this, one shows that there is a universal partial computable function $\text{Un}(k, x)$. This allows us to enumerate the partial computable functions. We will adopt the notation φ_k to denote the k -th unary partial computable function, defined by $\varphi_k(x) \simeq \text{Un}(k, x)$. (Kleene used $\{k\}$ for this purpose, but this notation has not been used as much recently.) Slightly more generally, we can uniformly enumerate the partial computable functions of arbitrary arities, and we will use φ_k^n to denote the k -th n -ary partial recursive function.

If $f(\vec{x}, y)$ is a total or partial function, then $\mu y f(\vec{x}, y)$ is the function of $\vec{x}$ that returns the least y such that $f(\vec{x}, y) = 0$, assuming that all of $f(\vec{x}, 0), \dots, f(\vec{x}, y-1)$ are defined; if there is no such y , $\mu y f(\vec{x}, y)$ is undefined. If $R(\vec{x}, y)$ is a relation, $\mu y R(\vec{x}, y)$ is defined to be the least y such that $R(\vec{x}, y)$ is true; in other words, the least y such that $1 \dot{-} \chi_R(\vec{x}, y) = 0$.

To show that a function is computable, there are two ways one can proceed:

1. Rigorously: describe a Turing machine or partial recursive function explicitly, and show that it computes the function you have in mind;
2. Informally: describe an algorithm that computes it, and appeal to Church's thesis.

There is no fine line between the two; a detailed description of an algorithm should provide enough information so that it is relatively clear how one could, in principle, design the right Turing machine or sequence of partial recursive definitions. Fully rigorous definitions are unlikely to be informative, and we will try to find a happy medium between these two approaches; in short, we will try to find intuitive yet rigorous proofs that the precise definitions could be obtained.

Créditos das Imagens

Referências Bibliográficas