

Capítulo udf

Tableaux

This chapter presents a signed analytic tableaux system.
To include or exclude material relevant to natural deduction as a proof system, use the “prfTab” tag.

-NoValue-

tab.1 Rules and Tableaux

fol:tab:rul:
sec A **tableau** is a systematic survey of the possible ways a **sentença** can be true or false in a **estrutura**. The building blocks of a tableau are **fórmulas assinadas**: **sentenças** plus a truth value “sign,” either $\mathbb{T}$ or $\mathbb{F}$. These signed **fórmulas** are arranged in a (downward growing) tree.

Definição tab.1. A **fórmula assinada** is a pair consisting of a truth value and a **sentença**, O.s., either:

$$\mathbb{T}\varphi \text{ or } \mathbb{F}\varphi.$$

Intuitively, we might read $\mathbb{T}\varphi$ as “ φ might be true” and $\mathbb{F}\varphi$ as “ φ might be false” (in some **estrutura**).

Each **fórmula assinada** in the tree is either an *assumption* (which are listed at the very top of the tree), or it is obtained from a **fórmula assinada** above it by one of a number of rules of inference. There are two rules for each possible **operador principal** of the preceding **fórmula**, one for the case where the sign is $\mathbb{T}$, and one for the case where the sign is $\mathbb{F}$. Some rules allow the tree to branch, and some only add **fórmulas assinadas** to the branch. A rule may be (and often must be) applied not to the immediately preceding **fórmula assinada**, but to any **fórmula assinada** in the branch from the root to the place the rule is applied.

A branch is *closed* when it contains both $\mathbb{T}\varphi$ and $\mathbb{F}\varphi$. A closed **tableau** is one where every branch is closed. Under the intuitive interpretation, any

branch describes a joint possibility, but $\mathbb{T}\varphi$ and $\mathbb{F}\varphi$ are not jointly possible. In other words, if a branch is closed, the possibility it describes has been ruled out. In particular, that means that a closed **tableau** rules out all possibilities of simultaneously making every assumption of the form $\mathbb{T}\varphi$ true and every assumption of the form $\mathbb{F}\varphi$ false.

A closed **tableau** for φ is a closed **tableau** with root $\mathbb{F}\varphi$. If such a closed **tableau** exists, all possibilities for φ being false have been ruled out; O.s., φ must be true in every **estrutur**.

-NoValue-

tab.2 Propositional Rules

Rules for $\neg$

fol:tab:prl:
sec

$$\frac{\mathbb{T}\neg\varphi}{\mathbb{F}\varphi} \neg\mathbb{T} \qquad \frac{\mathbb{F}\neg\varphi}{\mathbb{T}\varphi} \neg\mathbb{F}$$

Rules for $\wedge$

$$\frac{\mathbb{T}\varphi \wedge \psi}{\mathbb{T}\varphi \quad \mathbb{T}\psi} \wedge\mathbb{T} \qquad \frac{\mathbb{F}\varphi \wedge \psi}{\mathbb{F}\varphi \quad | \quad \mathbb{F}\psi} \wedge\mathbb{F}$$

Rules for $\vee$

$$\frac{\mathbb{T}\varphi \vee \psi}{\mathbb{T}\varphi \quad | \quad \mathbb{T}\psi} \vee\mathbb{T} \qquad \frac{\mathbb{F}\varphi \vee \psi}{\mathbb{F}\varphi \quad \mathbb{F}\psi} \vee\mathbb{F}$$

Rules for $\rightarrow$

$$\frac{\mathbb{T}\varphi \rightarrow \psi}{\mathbb{F}\varphi \quad | \quad \mathbb{T}\psi} \rightarrow\mathbb{T} \qquad \frac{\mathbb{F}\varphi \rightarrow \psi}{\mathbb{T}\varphi \quad \mathbb{F}\psi} \rightarrow\mathbb{F}$$

The Cut Rule

$$\frac{\overline{\mathbb{T} \varphi} \quad \overline{\mathbb{F} \varphi}}{\text{Cut}}$$

The Cut rule is not applied “to” a previous *fórmula assinada*; rather, it allows every branch in a *tableau* to be split in two, one branch containing $\mathbb{T} \varphi$, the other $\mathbb{F} \varphi$. It is not necessary—any set of *fórmulas assinadas* with a closed *tableau* has one not using Cut—but it allows us to combine *tableaux* in a convenient way.

-NoValue-

tab.3 Quantifier Rules

fol:tab:qrl:
sec Rules for $\forall$

$$\frac{\mathbb{T} \forall x \varphi(x)}{\mathbb{T} \varphi(t)} \forall \mathbb{T} \qquad \frac{\mathbb{F} \forall x \varphi(x)}{\mathbb{F} \varphi(a)} \forall \mathbb{F}$$

In $\forall \mathbb{T}$, t is a closed term (O.s., one without variables). In $\forall \mathbb{F}$, a is a *símbolo de constante* which must not occur anywhere in the branch above $\forall \mathbb{F}$ rule. We call a the *eigenvariable* of the $\forall \mathbb{F}$ inference.¹

Rules for $\exists$

$$\frac{\mathbb{T} \exists x \varphi(x)}{\mathbb{T} \varphi(a)} \exists \mathbb{T} \qquad \frac{\mathbb{F} \exists x \varphi(x)}{\mathbb{F} \varphi(t)} \exists \mathbb{F}$$

Again, t is a closed term, and a is a *símbolo de constante* which does not occur in the branch above the $\exists \mathbb{T}$ rule. We call a the *eigenvariable* of the $\exists \mathbb{T}$ inference.

The condition that an eigenvariable not occur in the branch above the $\forall \mathbb{F}$ or $\exists \mathbb{T}$ inference is called the *eigenvariable condition*.

Recall the convention that when φ is a *fórmula* with the *variável* x free, we indicate this by writing $\varphi(x)$. In the same context, $\varphi(t)$ then is short for $\varphi[t/x]$. So we could also write the $\exists \mathbb{F}$ rule as: explanation

¹We use the term “eigenvariable” even though a in the above rule is a *símbolo de constante*. This has historical reasons.

$$\frac{\mathbb{F} \exists x \varphi}{\mathbb{F} \varphi[t/x]} \exists \mathbb{F}$$

Note that t may already occur in φ , p.ex., φ might be $P(t, x)$. Thus, inferring $\mathbb{F} P(t, t)$ from $\mathbb{F} \exists x P(t, x)$ is a correct application of $\exists \mathbb{F}$. However, the eigenvariable conditions in $\forall \mathbb{F}$ and $\exists \mathbb{T}$ require that the **símbolo de constante** a does not occur in φ . So, you cannot correctly infer $\mathbb{F} P(a, a)$ from $\mathbb{F} \forall x P(a, x)$ using $\forall \mathbb{F}$.

explanation

In $\forall \mathbb{T}$ and $\exists \mathbb{F}$ there are no restrictions on the term t . On the other hand, in the $\exists \mathbb{T}$ and $\forall \mathbb{F}$ rules, the eigenvariable condition requires that the **símbolo de constante** a does not occur anywhere in the branches above the respective inference. It is necessary to ensure that the system is sound. Without this condition, the following would be a closed **tableau** for $\exists x \varphi(x) \rightarrow \forall x \varphi(x)$:

1.	$\mathbb{F} \exists x \varphi(x) \rightarrow \forall x \varphi(x)$	Assumption
2.	$\mathbb{T} \exists x \varphi(x)$	$\rightarrow \mathbb{F} 1$
3.	$\mathbb{F} \forall x \varphi(x)$	$\rightarrow \mathbb{F} 1$
4.	$\mathbb{T} \varphi(a)$	$\exists \mathbb{T} 2$
5.	$\mathbb{F} \varphi(a)$	$\forall \mathbb{F} 3$
	$\otimes$	

However, $\exists x \varphi(x) \rightarrow \forall x \varphi(x)$ is not valid.

-NoValue-

tab.4 Tableaux

explanation

We've said what an assumption is, and we've given the rules of inference. **Tableaux** are inductively generated from these: each **tableau** either is a single branch consisting of one or more assumptions, or it results from a **tableau** by applying one of the rules of inference on a branch.

fol:tab:der:
sec

Definição tab.2 (Tableau). A **tableau** for assumptions $S_1\varphi_1, \dots, S_n\varphi_n$ (where each S_i is either $\mathbb{T}$ or $\mathbb{F}$) is a finite tree of **fórmulas assinadas** satisfying the following conditions:

1. The n topmost **fórmulas assinadas** of the tree are $S_i\varphi_i$, one below the other.
2. Every **fórmula assinada** in the tree that is not one of the assumptions results from a correct application of an inference rule to a **fórmula assinada** in the branch above it.

A branch of a **tableau** is *closed* iff it contains both $\mathbb{T}\varphi$ and $\mathbb{F}\varphi$, and *open* otherwise. A **tableau** in which every branch is closed is a *closed tableau* (for its set of assumptions). If a **tableau** is not closed, O.s., if it contains at least one open branch, it is *open*.

Exemplo tab.3. Every set of assumptions on its own is a **tableau**, but it will generally not be closed. (Obviously, it is closed only if the assumptions already contain a pair of **fórmulas assinadas** $\mathbb{T}\varphi$ and $\mathbb{F}\varphi$.)

From a **tableau** (open or closed) we can obtain a new, larger one by applying one of the rules of inference to a **fórmula assinada** φ in it. The rule will append one or more **fórmulas assinadas** to the end of any branch containing the occurrence of φ to which we apply the rule.

For instance, consider the assumption $\mathbb{T}\varphi \wedge \neg\varphi$. Here is the (open) **tableau** consisting of just that assumption:

1. $\mathbb{T}\varphi \wedge \neg\varphi$ Assumption

We obtain a new **tableau** from it by applying the $\wedge\mathbb{T}$ rule to the assumption. That rule allows us to add two new lines to the **tableau**, $\mathbb{T}\varphi$ and $\mathbb{T}\neg\varphi$:

1. $\mathbb{T}\varphi \wedge \neg\varphi$ Assumption
2. $\mathbb{T}\varphi$ $\wedge\mathbb{T} 1$
3. $\mathbb{T}\neg\varphi$ $\wedge\mathbb{T} 1$

When we write down **tableaux**, we record the rules we've applied on the right (p.ex., $\wedge\mathbb{T}1$ means that the **fórmula assinada** on that line is the result of applying the $\wedge\mathbb{T}$ rule to the **fórmula assinada** on line 1). This new **tableau** now contains additional **fórmulas assinadas**, but to only one ($\mathbb{T}\neg\varphi$) can we apply a rule (in this case, the $\neg\mathbb{T}$ rule). This results in the closed **tableau**

1. $\mathbb{T}\varphi \wedge \neg\varphi$ Assumption
 2. $\mathbb{T}\varphi$ $\wedge\mathbb{T} 1$
 3. $\mathbb{T}\neg\varphi$ $\wedge\mathbb{T} 1$
 4. $\mathbb{F}\varphi$ $\neg\mathbb{T} 3$
- $\otimes$

-NoValue-

tab.5 Examples of Tableaux

fol:tab:pro:
sec

Exemplo tab.4. Let's find a closed **tableau** for the **sentença** $(\varphi \wedge \psi) \rightarrow \varphi$.

We begin by writing the corresponding assumption at the top of the **tableau**.

1. $\mathbb{F}(\varphi \wedge \psi) \rightarrow \varphi$ Assumption

There is only one assumption, so only one **fórmula assinada** to which we can apply a rule. (For every **fórmula assinada**, there is always at most one rule that can be applied: it's the rule for the corresponding sign and **operador principal** of the **sentença**.) In this case, this means, we must apply $\rightarrow\mathbb{F}$.

1.	$\mathbb{F}(\varphi \wedge \psi) \rightarrow \varphi \checkmark$	Assumption
2.	$\mathbb{T}\varphi \wedge \psi$	$\rightarrow\mathbb{F} 1$
3.	$\mathbb{F}\varphi$	$\rightarrow\mathbb{F} 1$

To keep track of which **fórmulas assinadas** we have applied their corresponding rules to, we write a checkmark next to the sentence. However, *only* write a checkmark if the rule has been applied to all open branches. Once a **fórmula assinada** has had the corresponding rule applied in every open branch, we will not have to return to it and apply the rule again. In this case, there is only one branch, so the rule only has to be applied once. (Note that checkmarks are only a convenience for constructing tableaux and are not officially part of the syntax of tableaux.)

There is one new **fórmula assinada** to which we can apply a rule: the $\mathbb{T}\varphi \wedge \psi$ on line 2. Applying the $\wedge\mathbb{T}$ rule results in:

1.	$\mathbb{F}(\varphi \wedge \psi) \rightarrow \varphi \checkmark$	Assumption
2.	$\mathbb{T}\varphi \wedge \psi \checkmark$	$\rightarrow\mathbb{F} 1$
3.	$\mathbb{F}\varphi$	$\rightarrow\mathbb{F} 1$
4.	$\mathbb{T}\varphi$	$\wedge\mathbb{T} 2$
5.	$\mathbb{T}\psi$	$\wedge\mathbb{T} 2$
	$\otimes$	

Since the branch now contains both $\mathbb{T}\varphi$ (on line 4) and $\mathbb{F}\varphi$ (on line 3), the branch is closed. Since it is the only branch, the **tableau** is closed. We have found a closed **tableau** for $(\varphi \wedge \psi) \rightarrow \varphi$.

Exemplo tab.5. Now let's find a closed **tableau** for $(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)$.

We begin with the corresponding assumption:

1.	$\mathbb{F}(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)$	Assumption
----	--	------------

The one **fórmula assinada** in this **tableau** has **operador principal** $\rightarrow$ and sign $\mathbb{F}$, so we apply the $\rightarrow\mathbb{F}$ rule to it to obtain:

1.	$\mathbb{F}(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi) \checkmark$	Assumption
2.	$\mathbb{T}\neg\varphi \vee \psi$	$\rightarrow\mathbb{F} 1$
3.	$\mathbb{F}(\varphi \rightarrow \psi)$	$\rightarrow\mathbb{F} 1$

We now have a choice as to whether to apply $\vee\mathbb{T}$ to line 2 or $\rightarrow\mathbb{F}$ to line 3. It actually doesn't matter which order we pick, as long as each **fórmula assinada** has its corresponding rule applied in every branch. So let's pick the first one. The $\vee\mathbb{T}$ rule allows the **tableau** to branch, and the two conclusions of the rule will be the new **fórmulas assinadas** added to the two new branches. This results in:

1.	$\mathbb{F}(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi) \checkmark$	Assumption
2.	$\mathbb{T}\neg\varphi \vee \psi \checkmark$	$\rightarrow\mathbb{F} 1$
3.	$\mathbb{F}(\varphi \rightarrow \psi)$	$\rightarrow\mathbb{F} 1$
$\swarrow \quad \searrow$		
4.	$\mathbb{T}\neg\varphi \quad \mathbb{T}\psi$	$\vee\mathbb{T} 2$

We have not applied the $\rightarrow\mathbb{F}$ rule to line 3 yet: let's do that now. To save time, we apply it to both branches. Recall that we write a checkmark next to a **fórmula assinada** only if we have applied the corresponding rule in every open branch. So it's a good idea to apply a rule at the end of every branch that contains the **fórmula assinada** the rule applies to. That way we won't have to return to that **fórmula assinada** lower down in the various branches.

1.	$\mathbb{F}(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi) \checkmark$	Assumption
2.	$\mathbb{T}\neg\varphi \vee \psi \checkmark$	$\rightarrow\mathbb{F} 1$
3.	$\mathbb{F}(\varphi \rightarrow \psi) \checkmark$	$\rightarrow\mathbb{F} 1$
$\swarrow \quad \searrow$		
4.	$\mathbb{T}\neg\varphi \quad \mathbb{T}\psi$	$\vee\mathbb{T} 2$
5.	$\mathbb{T}\varphi \quad \mathbb{T}\varphi$	$\rightarrow\mathbb{F} 3$
6.	$\mathbb{F}\psi \quad \mathbb{F}\psi$	$\rightarrow\mathbb{F} 3$
$\quad \quad \quad \otimes$		

The right branch is now closed. On the left branch, we can still apply the $\neg\mathbb{T}$ rule to line 4. This results in $\mathbb{F}\varphi$ and closes the left branch:

1.	$\mathbb{F}(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi) \checkmark$	Assumption
2.	$\mathbb{T}\neg\varphi \vee \psi \checkmark$	$\rightarrow\mathbb{F} 1$
3.	$\mathbb{F}(\varphi \rightarrow \psi) \checkmark$	$\rightarrow\mathbb{F} 1$
$\swarrow \quad \searrow$		
4.	$\mathbb{T}\neg\varphi \quad \mathbb{T}\psi$	$\vee\mathbb{T} 2$
5.	$\mathbb{T}\varphi \quad \mathbb{T}\varphi$	$\rightarrow\mathbb{F} 3$
6.	$\mathbb{F}\psi \quad \mathbb{F}\psi$	$\rightarrow\mathbb{F} 3$
7.	$\mathbb{F}\varphi \quad \otimes$	$\neg\mathbb{T} 4$
$\quad \quad \quad \otimes$		

Exemplo tab.6. We can give **tableaux** for any number of **fórmulas assinadas** as assumptions. Often it is also necessary to apply more than one rule that allows branching; and in general a **tableau** can have any number of branches. For instance, consider a **tableau** for $\{\mathbb{T}\varphi \vee (\psi \wedge \chi), \mathbb{F}(\varphi \vee \psi) \wedge (\varphi \vee \chi)\}$. We start by applying the $\vee\mathbb{T}$ to the first assumption:

1.	$\mathbb{T}\varphi \vee (\psi \wedge \chi) \checkmark$	Assumption
2.	$\mathbb{F}(\varphi \vee \psi) \wedge (\varphi \vee \chi)$	Assumption
$\swarrow \quad \searrow$		
3.	$\mathbb{T}\varphi \quad \mathbb{T}\psi \wedge \chi$	$\vee\mathbb{T} 1$

Now we can apply the $\wedge\mathbb{F}$ rule to line 2. We do this on both branches simultaneously, and can therefore check off line 2:

1.	$\mathbb{T}\varphi \vee (\psi \wedge \chi) \checkmark$	Assumption
2.	$\mathbb{F}(\varphi \vee \psi) \wedge (\varphi \vee \chi) \checkmark$	Assumption
3.	$\begin{array}{cc} \mathbb{T}\varphi & \mathbb{T}\psi \wedge \chi \end{array}$	$\vee\mathbb{T} 1$
4.	$\begin{array}{cc} \mathbb{F}\varphi \vee \psi & \mathbb{F}\varphi \vee \chi \end{array} \quad \begin{array}{cc} \mathbb{F}\varphi \vee \psi & \mathbb{F}\varphi \vee \chi \end{array}$	$\wedge\mathbb{F} 2$

Now we can apply $\vee\mathbb{F}$ to all the branches containing $\varphi \vee \psi$:

1.	$\mathbb{T}\varphi \vee (\psi \wedge \chi) \checkmark$	Assumption
2.	$\mathbb{F}(\varphi \vee \psi) \wedge (\varphi \vee \chi) \checkmark$	Assumption
3.	$\begin{array}{cc} \mathbb{T}\varphi & \mathbb{T}\psi \wedge \chi \end{array}$	$\vee\mathbb{T} 1$
4.	$\begin{array}{cc} \mathbb{F}\varphi \vee \psi \checkmark & \mathbb{F}\varphi \vee \chi \end{array} \quad \begin{array}{cc} \mathbb{F}\varphi \vee \psi \checkmark & \mathbb{F}\varphi \vee \chi \end{array}$	$\wedge\mathbb{F} 2$
5.	$\begin{array}{cc} \mathbb{F}\varphi & \mathbb{F}\varphi \end{array}$	$\vee\mathbb{F} 4$
6.	$\begin{array}{cc} \mathbb{F}\psi & \mathbb{F}\psi \end{array}$	$\vee\mathbb{F} 4$
	$\otimes$	

The leftmost branch is now closed. Let's now apply $\vee\mathbb{F}$ to $\varphi \vee \chi$:

1.	$\mathbb{T}\varphi \vee (\psi \wedge \chi) \checkmark$	Assumption
2.	$\mathbb{F}(\varphi \vee \psi) \wedge (\varphi \vee \chi) \checkmark$	Assumption
3.	$\begin{array}{cc} \mathbb{T}\varphi & \mathbb{T}\psi \wedge \chi \end{array}$	$\vee\mathbb{T} 1$
4.	$\begin{array}{cc} \mathbb{F}\varphi \vee \psi \checkmark & \mathbb{F}\varphi \vee \chi \checkmark \end{array} \quad \begin{array}{cc} \mathbb{F}\varphi \vee \psi \checkmark & \mathbb{F}\varphi \vee \chi \checkmark \end{array}$	$\wedge\mathbb{F} 2$
5.	$\begin{array}{cc} \mathbb{F}\varphi & \mathbb{F}\varphi \end{array}$	$\vee\mathbb{F} 4$
6.	$\begin{array}{cc} \mathbb{F}\psi & \mathbb{F}\psi \end{array}$	$\vee\mathbb{F} 4$
7.	$\begin{array}{cc} \otimes & \mathbb{F}\varphi \end{array}$	$\vee\mathbb{F} 4$
8.	$\begin{array}{cc} & \mathbb{F}\chi \end{array}$	$\vee\mathbb{F} 4$
	$\otimes$	

Note that we moved the result of applying $\vee\mathbb{F}$ a second time below for clarity. In this instance it would not have been needed, since the justifications would have been the same.

Two branches remain open, and $\mathbb{T}\psi \wedge \chi$ on line 3 remains unchecked. We apply $\wedge\mathbb{T}$ to it to obtain a closed **tableau**:

1.	$\mathbb{T} \varphi \vee (\psi \wedge \chi) \checkmark$	Assumption
2.	$\mathbb{F} (\varphi \vee \psi) \wedge (\varphi \vee \chi) \checkmark$	Assumption
3.	$\mathbb{T} \varphi$ $\mathbb{T} \psi \wedge \chi \checkmark$	$\vee \mathbb{T} 1$
4.	$\mathbb{F} \varphi \vee \psi \checkmark$ $\mathbb{F} \varphi \vee \chi \checkmark$ $\mathbb{F} \varphi \vee \psi \checkmark$ $\mathbb{F} \varphi \vee \chi \checkmark$	$\wedge \mathbb{F} 2$
5.	$\mathbb{F} \varphi$ $\mathbb{F} \varphi$ $\mathbb{F} \varphi$ $\mathbb{F} \varphi$	$\vee \mathbb{F} 4$
6.	$\mathbb{F} \psi$ $\mathbb{F} \chi$ $\mathbb{F} \psi$ $\mathbb{F} \chi$	$\vee \mathbb{F} 4$
7.	$\otimes$ $\otimes$ $\mathbb{T} \psi$ $\mathbb{T} \psi$	$\wedge \mathbb{T} 3$
8.	$\otimes$ $\otimes$ $\mathbb{T} \chi$ $\mathbb{T} \chi$	$\wedge \mathbb{T} 3$
	$\otimes$ $\otimes$	

For comparison, here's a closed **tableau** for the same set of assumptions in which the rules are applied in a different order:

1.	$\mathbb{T} \varphi \vee (\psi \wedge \chi) \checkmark$	Assumption
2.	$\mathbb{F} (\varphi \vee \psi) \wedge (\varphi \vee \chi) \checkmark$	Assumption
3.	$\mathbb{F} \varphi \vee \psi \checkmark$ $\mathbb{F} \varphi \vee \chi \checkmark$	$\wedge \mathbb{F} 2$
4.	$\mathbb{F} \varphi$ $\mathbb{F} \varphi$	$\vee \mathbb{F} 3$
5.	$\mathbb{F} \psi$ $\mathbb{F} \chi$	$\vee \mathbb{F} 3$
6.	$\mathbb{T} \varphi$ $\mathbb{T} \psi \wedge \chi \checkmark$ $\mathbb{T} \varphi$ $\mathbb{T} \psi \wedge \chi \checkmark$	$\vee \mathbb{T} 1$
7.	$\otimes$ $\mathbb{T} \psi$ $\otimes$ $\mathbb{T} \psi$	$\wedge \mathbb{T} 6$
8.	$\otimes$ $\mathbb{T} \chi$ $\otimes$ $\mathbb{T} \chi$	$\wedge \mathbb{T} 6$
	$\otimes$ $\otimes$	

Problema tab.1. Give closed **tableaux** of the following:

1. $\mathbb{T} \varphi \wedge (\psi \wedge \chi), \mathbb{F} (\varphi \wedge \psi) \wedge \chi$.
2. $\mathbb{T} \varphi \vee (\psi \vee \chi), \mathbb{F} (\varphi \vee \psi) \vee \chi$.
3. $\mathbb{T} \varphi \rightarrow (\psi \rightarrow \chi), \mathbb{F} \psi \rightarrow (\varphi \rightarrow \chi)$.
4. $\mathbb{T} \varphi, \mathbb{F} \neg \neg \varphi$.

Problema tab.2. Give closed **tableaux** of the following:

1. $\mathbb{T} (\varphi \vee \psi) \rightarrow \chi, \mathbb{F} \varphi \rightarrow \chi$.
2. $\mathbb{T} (\varphi \rightarrow \chi) \wedge (\psi \rightarrow \chi), \mathbb{F} (\varphi \vee \psi) \rightarrow \chi$.
3. $\mathbb{F} \neg (\varphi \wedge \neg \varphi)$.
4. $\mathbb{T} \psi \rightarrow \varphi, \mathbb{F} \neg \varphi \rightarrow \neg \psi$.

5. $\mathbb{F} (\varphi \rightarrow \neg\varphi) \rightarrow \neg\varphi.$
6. $\mathbb{F} \neg(\varphi \rightarrow \psi) \rightarrow \neg\psi.$
7. $\mathbb{T} \varphi \rightarrow \chi, \mathbb{F} \neg(\varphi \wedge \neg\chi).$
8. $\mathbb{T} \varphi \wedge \neg\chi, \mathbb{F} \neg(\varphi \rightarrow \chi).$
9. $\mathbb{T} \varphi \vee \psi, \neg\psi, \mathbb{F} \varphi.$
10. $\mathbb{T} \neg\varphi \vee \neg\psi, \mathbb{F} \neg(\varphi \wedge \psi).$
11. $\mathbb{F} (\neg\varphi \wedge \neg\psi) \rightarrow \neg(\varphi \vee \psi).$
12. $\mathbb{F} \neg(\varphi \vee \psi) \rightarrow (\neg\varphi \wedge \neg\psi).$

Problema tab.3. Give closed **tableaux** of the following:

1. $\mathbb{T} \neg(\varphi \rightarrow \psi), \mathbb{F} \varphi.$
2. $\mathbb{T} \neg(\varphi \wedge \psi), \mathbb{F} \neg\varphi \vee \neg\psi.$
3. $\mathbb{T} \varphi \rightarrow \psi, \mathbb{F} \neg\varphi \vee \psi.$
4. $\mathbb{F} \neg\neg\varphi \rightarrow \varphi.$
5. $\mathbb{T} \varphi \rightarrow \psi, \mathbb{T} \neg\varphi \rightarrow \psi, \mathbb{F} \psi.$
6. $\mathbb{T} (\varphi \wedge \psi) \rightarrow \chi, \mathbb{F} (\varphi \rightarrow \chi) \vee (\psi \rightarrow \chi).$
7. $\mathbb{T} (\varphi \rightarrow \psi) \rightarrow \varphi, \mathbb{F} \varphi.$
8. $\mathbb{F} (\varphi \rightarrow \psi) \vee (\psi \rightarrow \chi).$

-NoValue-

tab.6 Tableaux with Quantifiers

Exemplo tab.7. When dealing with quantifiers, we have to make sure not to violate the eigenvariable condition, and sometimes this requires us to play around with the order of carrying out certain inferences. In general, it helps to try and take care of rules subject to the eigenvariable condition first (they will be higher up in the finished **tableau**).

Let's see how we'd give a **tableau** for the **sentença** $\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)$. Starting as usual, we start by recording the assumption,

$$1. \quad \mathbb{F} \exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x) \quad \text{Assumption}$$

Since the **operador principal** is $\rightarrow$, we apply the $\rightarrow\mathbb{F}$:

1.	$\mathbb{F} \exists x \neg \varphi(x) \rightarrow \neg \forall x \varphi(x) \checkmark$	Assumption
2.	$\mathbb{T} \exists x \neg \varphi(x)$	$\rightarrow \mathbb{F} 1$
3.	$\mathbb{F} \neg \forall x \varphi(x)$	$\rightarrow \mathbb{F} 1$

The next line to deal with is 2. We use $\exists\mathbb{T}$. This requires a new **símbolo de constante**; since no **símbolos de constante** yet occur, we can pick any one, say, a .

1.	$\mathbb{F} \exists x \neg \varphi(x) \rightarrow \neg \forall x \varphi(x) \checkmark$	Assumption
2.	$\mathbb{T} \exists x \neg \varphi(x) \checkmark$	$\rightarrow \mathbb{F} 1$
3.	$\mathbb{F} \neg \forall x \varphi(x)$	$\rightarrow \mathbb{F} 1$
4.	$\mathbb{T} \neg \varphi(a)$	$\exists\mathbb{T} 2$

Now we apply $\neg\mathbb{F}$ to line 3:

1.	$\mathbb{F} \exists x \neg \varphi(x) \rightarrow \neg \forall x \varphi(x) \checkmark$	Assumption
2.	$\mathbb{T} \exists x \neg \varphi(x) \checkmark$	$\rightarrow \mathbb{F} 1$
3.	$\mathbb{F} \neg \forall x \varphi(x) \checkmark$	$\rightarrow \mathbb{F} 1$
4.	$\mathbb{T} \neg \varphi(a)$	$\exists\mathbb{T} 2$
5.	$\mathbb{T} \forall x \varphi(x)$	$\neg\mathbb{F} 3$

We obtain a closed **tableau** by applying $\neg\mathbb{T}$ to line 4, followed by $\forall\mathbb{T}$ to line 5.

1.	$\mathbb{F} \exists x \neg \varphi(x) \rightarrow \neg \forall x \varphi(x) \checkmark$	Assumption
2.	$\mathbb{T} \exists x \neg \varphi(x) \checkmark$	$\rightarrow \mathbb{F} 1$
3.	$\mathbb{F} \neg \forall x \varphi(x) \checkmark$	$\rightarrow \mathbb{F} 1$
4.	$\mathbb{T} \neg \varphi(a)$	$\exists\mathbb{T} 2$
5.	$\mathbb{T} \forall x \varphi(x)$	$\neg\mathbb{F} 3$
6.	$\mathbb{F} \varphi(a)$	$\neg\mathbb{T} 4$
7.	$\mathbb{T} \varphi(a)$	$\forall\mathbb{T} 5$
	$\otimes$	

Exemplo tab.8. Let's see how we'd give a **tableau** for the set

$$\mathbb{F} \exists x \chi(x, b), \mathbb{T} \exists x (\varphi(x) \wedge \psi(x)), \mathbb{T} \forall x (\psi(x) \rightarrow \chi(x, b)).$$

Starting as usual, we start with the assumptions:

1.	$\mathbb{F} \exists x \chi(x, b)$	Assumption
2.	$\mathbb{T} \exists x (\varphi(x) \wedge \psi(x))$	Assumption
3.	$\mathbb{T} \forall x (\psi(x) \rightarrow \chi(x, b))$	Assumption

We should always apply a rule with the eigenvariable condition first; in this case that would be $\exists\mathbb{T}$ to line 2. Since the assumptions contain the **símbolo de constante** b , we have to use a different one; let's pick a again.

1.	$\mathbb{F} \exists x \chi(x, b)$	Assumption
2.	$\mathbb{T} \exists x (\varphi(x) \wedge \psi(x)) \checkmark$	Assumption
3.	$\mathbb{T} \forall x (\psi(x) \rightarrow \chi(x, b))$	Assumption
4.	$\mathbb{T} \varphi(a) \wedge \psi(a)$	$\exists\mathbb{T} 2$

If we now apply $\exists\mathbb{F}$ to line 1 or $\forall\mathbb{T}$ to line 3, we have to decide which term t to substitute for x . Since there is no eigenvariable condition for these rules, we can pick any term we like. In some cases we may even have to apply the rule several times with different ts . But as a general rule, it pays to pick one of the terms already occurring in the **tableau**—in this case, a and b —and in this case we can guess that a will be more likely to result in a closed branch.

1.	$\mathbb{F} \exists x \chi(x, b)$	Assumption
2.	$\mathbb{T} \exists x (\varphi(x) \wedge \psi(x)) \checkmark$	Assumption
3.	$\mathbb{T} \forall x (\psi(x) \rightarrow \chi(x, b))$	Assumption
4.	$\mathbb{T} \varphi(a) \wedge \psi(a)$	$\exists\mathbb{T} 2$
5.	$\mathbb{F} \chi(a, b)$	$\exists\mathbb{F} 1$
6.	$\mathbb{T} \psi(a) \rightarrow \chi(a, b)$	$\forall\mathbb{T} 3$

We don't check the **fórmulas assinadas** in lines 1 and 3, since we may have to use them again. Now apply $\wedge\mathbb{T}$ to line 4:

1.	$\mathbb{F} \exists x \chi(x, b)$	Assumption
2.	$\mathbb{T} \exists x (\varphi(x) \wedge \psi(x)) \checkmark$	Assumption
3.	$\mathbb{T} \forall x (\psi(x) \rightarrow \chi(x, b))$	Assumption
4.	$\mathbb{T} \varphi(a) \wedge \psi(a) \checkmark$	$\exists\mathbb{T} 2$
5.	$\mathbb{F} \chi(a, b)$	$\exists\mathbb{F} 1$
6.	$\mathbb{T} \psi(a) \rightarrow \chi(a, b)$	$\forall\mathbb{T} 3$
7.	$\mathbb{T} \varphi(a)$	$\wedge\mathbb{T} 4$
8.	$\mathbb{T} \psi(a)$	$\wedge\mathbb{T} 4$

If we now apply $\rightarrow\mathbb{T}$ to line 6, the **tableau** closes:

1.	$\mathbb{F} \exists x \chi(x, b)$	Assumption
2.	$\mathbb{T} \exists x (\varphi(x) \wedge \psi(x)) \checkmark$	Assumption
3.	$\mathbb{T} \forall x (\psi(x) \rightarrow \chi(x, b))$	Assumption
4.	$\mathbb{T} \varphi(a) \wedge \psi(a) \checkmark$	$\exists\mathbb{T} 2$
5.	$\mathbb{F} \chi(a, b)$	$\exists\mathbb{F} 1$
6.	$\mathbb{T} \psi(a) \rightarrow \chi(a, b) \checkmark$	$\forall\mathbb{T} 3$
7.	$\mathbb{T} \varphi(a)$	$\wedge\mathbb{T} 4$
8.	$\mathbb{T} \psi(a)$	$\wedge\mathbb{T} 4$
9.	$\begin{array}{c} \swarrow \quad \searrow \\ \mathbb{F} \psi(a) \quad \mathbb{T} \chi(a, b) \\ \otimes \quad \quad \otimes \end{array}$	$\rightarrow\mathbb{T} 6$

Exemplo tab.9. We construct a **tableau** for the set

$$\mathbb{T} \forall x \varphi(x), \mathbb{T} \forall x \varphi(x) \rightarrow \exists y \psi(y), \mathbb{T} \neg \exists y \psi(y).$$

Starting as usual, we write down the assumptions:

- | | | |
|----|---|------------|
| 1. | $\mathbb{T} \forall x \varphi(x)$ | Assumption |
| 2. | $\mathbb{T} \forall x \varphi(x) \rightarrow \exists y \psi(y)$ | Assumption |
| 3. | $\mathbb{T} \neg \exists y \psi(y)$ | Assumption |

We begin by applying the $\neg\mathbb{T}$ rule to line 3. A corollary to the rule “always apply rules with eigenvariable conditions first” is “defer applying quantifier rules without eigenvariable conditions until needed.” Also, defer rules that result in a split.

- | | | |
|----|---|--------------------|
| 1. | $\mathbb{T} \forall x \varphi(x)$ | Assumption |
| 2. | $\mathbb{T} \forall x \varphi(x) \rightarrow \exists y \psi(y)$ | Assumption |
| 3. | $\mathbb{T} \neg \exists y \psi(y) \checkmark$ | Assumption |
| 4. | $\mathbb{F} \exists y \psi(y)$ | $\neg\mathbb{T} 3$ |

The new line 4 requires $\exists\mathbb{F}$, a quantifier rule without the eigenvariable condition. So we defer this in favor of using $\rightarrow\mathbb{T}$ on line 2.

- | | | |
|---------------------------|--|---------------------------|
| 1. | $\mathbb{T} \forall x \varphi(x)$ | Assumption |
| 2. | $\mathbb{T} \forall x \varphi(x) \rightarrow \exists y \psi(y) \checkmark$ | Assumption |
| 3. | $\mathbb{T} \neg \exists y \psi(y) \checkmark$ | Assumption |
| 4. | $\mathbb{F} \exists y \psi(y)$ | $\neg\mathbb{T} 3$ |
| $\swarrow \quad \searrow$ | | |
| 5. | $\mathbb{F} \forall x \varphi(x) \quad \mathbb{T} \exists y \psi(y)$ | $\rightarrow\mathbb{T} 2$ |

Both new **fórmulas assinadas** require rules with eigenvariable conditions, so these should be next:

- | | | |
|---------------------------|--|--|
| 1. | $\mathbb{T} \forall x \varphi(x)$ | Assumption |
| 2. | $\mathbb{T} \forall x \varphi(x) \rightarrow \exists y \psi(y) \checkmark$ | Assumption |
| 3. | $\mathbb{T} \neg \exists y \psi(y) \checkmark$ | Assumption |
| 4. | $\mathbb{F} \exists y \psi(y)$ | $\neg\mathbb{T} 3$ |
| $\swarrow \quad \searrow$ | | |
| 5. | $\mathbb{F} \forall x \varphi(x) \checkmark \quad \mathbb{T} \exists y \psi(y) \checkmark$ | $\rightarrow\mathbb{T} 2$ |
| 6. | $\mathbb{F} \varphi(b) \quad \mathbb{T} \psi(c)$ | $\forall\mathbb{F} 5; \exists\mathbb{T} 5$ |

To close the branches, we have to use the **fórmulas assinadas** on lines 1 and 3. The corresponding rules ($\forall\mathbb{T}$ and $\exists\mathbb{F}$) don't have eigenvariable conditions, so we are free to pick whichever terms are suitable. In this case, that's b and c , respectively.

1.	$\mathbb{T} \forall x \varphi(x)$	Assumption	
2.	$\mathbb{T} \forall x \varphi(x) \rightarrow \exists y \psi(y) \checkmark$	Assumption	
3.	$\mathbb{T} \neg \exists y \psi(y) \checkmark$	Assumption	
4.	$\mathbb{F} \exists y \psi(y)$	$\neg \mathbb{T} 3$	
$\swarrow \qquad \searrow$			
5.	$\mathbb{F} \forall x \varphi(x) \checkmark$	$\mathbb{T} \exists y \psi(y) \checkmark$	$\rightarrow \mathbb{T} 2$
6.	$\mathbb{F} \varphi(b)$	$\mathbb{T} \psi(c)$	$\forall \mathbb{F} 5; \exists \mathbb{T} 5$
7.	$\mathbb{T} \varphi(b)$	$\mathbb{F} \psi(c)$	$\forall \mathbb{T} 1; \exists \mathbb{F} 4$
	$\otimes$	$\otimes$	

Problema tab.4. Give closed **tableaux** of the following:

1. $\mathbb{F} (\forall x \varphi(x) \wedge \forall y \psi(y)) \rightarrow \forall z (\varphi(z) \wedge \psi(z))$.
2. $\mathbb{F} (\exists x \varphi(x) \vee \exists y \psi(y)) \rightarrow \exists z (\varphi(z) \vee \psi(z))$.
3. $\mathbb{T} \forall x (\varphi(x) \rightarrow \psi), \mathbb{F} \exists y \varphi(y) \rightarrow \psi$.
4. $\mathbb{T} \forall x \neg \varphi(x), \mathbb{F} \neg \exists x \varphi(x)$.
5. $\mathbb{F} \neg \exists x \varphi(x) \rightarrow \forall x \neg \varphi(x)$.
6. $\mathbb{F} \neg \exists x \forall y ((\varphi(x, y) \rightarrow \neg \varphi(y, y)) \wedge (\neg \varphi(y, y) \rightarrow \varphi(x, y)))$.

Problema tab.5. Give closed **tableaux** of the following:

1. $\mathbb{F} \neg \forall x \varphi(x) \rightarrow \exists x \neg \varphi(x)$.
2. $\mathbb{T} (\forall x \varphi(x) \rightarrow \psi), \mathbb{F} \exists y (\varphi(y) \rightarrow \psi)$.
3. $\mathbb{F} \exists x (\varphi(x) \rightarrow \forall y \varphi(y))$.

-NoValue-

tab.7 Proof-Theoretic Notions

fol:tab:ptn:
sec

This section collects the definitions of the provability relation and consistency for tableaux.

explanation Just as we've defined a number of important semantic notions (validity, entailment, satisfiability), we now define corresponding *proof-theoretic notions*. These are not defined by appeal to satisfaction of **sentenças** in **estruturas**, but by appeal to the existence of certain closed **tableaux**. It was an important discovery that these notions coincide. That they do is the content of the *soundness* and *completeness theorems*.

Definição tab.10 (Theorems). A *sentença* φ is a *theorem* if there is a closed *tableau* for $\mathbb{F} \varphi$. We write $\vdash \varphi$ if φ is a theorem and $\nvdash \varphi$ if it is not.

Definição tab.11 (Derivabilidade). A *sentença* φ is *derivável* from a set of *sentenças* Γ , $\Gamma \vdash \varphi$ iff there is a finite set $\{\psi_1, \dots, \psi_n\} \subseteq \Gamma$ and a closed *tableau* for the set

$$\{\mathbb{F} \varphi, \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}.$$

If φ is not *derivável* from Γ we write $\Gamma \nvdash \varphi$.

Definição tab.12 (Consistency). A set of *sentenças* Γ is *inconsistent* iff there is a finite set $\{\psi_1, \dots, \psi_n\} \subseteq \Gamma$ and a closed *tableau* for the set

$$\{\mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}.$$

If Γ is not inconsistent, we say it is *consistent*.

fol:tab:ptn:
prop:reflexivity

Proposição tab.13 (Reflexivity). If $\varphi \in \Gamma$, then $\Gamma \vdash \varphi$.

Prova. If $\varphi \in \Gamma$, $\{\varphi\}$ is a finite subset of Γ and the *tableau*

1. $\mathbb{F} \varphi$ Assumption
2. $\mathbb{T} \varphi$ Assumption
- $\otimes$

is closed. □

fol:tab:ptn:
prop:monotonicity

Proposição tab.14 (Monotonicity). If $\Gamma \subseteq \Delta$ and $\Gamma \vdash \varphi$, then $\Delta \vdash \varphi$.

Prova. Any finite subset of Γ is also a finite subset of Δ . □

fol:tab:ptn:
prop:transitivity

Proposição tab.15 (Transitivity). If $\Gamma \vdash \varphi$ and $\{\varphi\} \cup \Delta \vdash \psi$, then $\Gamma \cup \Delta \vdash \psi$.

Prova. If $\{\varphi\} \cup \Delta \vdash \psi$, then there is a finite subset $\Delta_0 = \{\chi_1, \dots, \chi_n\} \subseteq \Delta$ such that

$$\{\mathbb{F} \psi, \mathbb{T} \varphi, \mathbb{T} \chi_1, \dots, \mathbb{T} \chi_n\}$$

has a closed *tableau*. If $\Gamma \vdash \varphi$ then there are $\theta_1, \dots, \theta_m$ such that

$$\{\mathbb{F} \varphi, \mathbb{T} \theta_1, \dots, \mathbb{T} \theta_m\}$$

has a closed *tableau*.

Now consider the *tableau* with assumptions

$$\mathbb{F} \psi, \mathbb{T} \chi_1, \dots, \mathbb{T} \chi_n, \mathbb{T} \theta_1, \dots, \mathbb{T} \theta_m.$$

Apply the Cut rule on φ . This generates two branches, one has $\mathbb{T}\varphi$ in it, the other $\mathbb{F}\varphi$. Thus, on the one branch, all of

$$\{\mathbb{F}\psi, \mathbb{T}\varphi, \mathbb{T}\chi_1, \dots, \mathbb{T}\chi_n\}$$

are available. Since there is a closed **tableau** for these assumptions, we can attach it to that branch; every branch through $\mathbb{T}\varphi$ closes. On the other branch, all of

$$\{\mathbb{F}\varphi, \mathbb{T}\theta_1, \dots, \mathbb{T}\theta_m\}$$

are available, so we can also complete the other side to obtain a closed **tableau**. This shows $\Gamma \cup \Delta \vdash \psi$. $\square$

Note that this means that in particular if $\Gamma \vdash \varphi$ and $\varphi \vdash \psi$, then $\Gamma \vdash \psi$. It follows also that if $\varphi_1, \dots, \varphi_n \vdash \psi$ and $\Gamma \vdash \varphi_i$ for each i , then $\Gamma \vdash \psi$.

Proposição tab.16. Γ is inconsistent iff $\Gamma \vdash \varphi$ for every *sentença* φ .

*fol:tab:ptn:
prop:incons*

Prova. Exercise. $\square$

Problema tab.6. Prove **Proposição tab.16**

Proposição tab.17 (Compactness).

*fol:tab:ptn:
prop:proves-compact*

1. If $\Gamma \vdash \varphi$ then there is a finite subset $\Gamma_0 \subseteq \Gamma$ such that $\Gamma_0 \vdash \varphi$.
2. If every finite subset of Γ is consistent, then Γ is consistent.

Prova. 1. If $\Gamma \vdash \varphi$, then there is a finite subset $\Gamma_0 = \{\psi_1, \dots, \psi_n\}$ and a closed **tableau** for

$$\{\mathbb{F}\varphi, \mathbb{T}\psi_1, \dots, \mathbb{T}\psi_n\}$$

This **tableau** also shows $\Gamma_0 \vdash \varphi$.

2. If Γ is inconsistent, then for some finite subset $\Gamma_0 = \{\psi_1, \dots, \psi_n\}$ there is a closed **tableau** for

$$\{\mathbb{T}\psi_1, \dots, \mathbb{T}\psi_n\}$$

This closed **tableau** shows that Γ_0 is inconsistent. $\square$

-NoValue-

tab.8 Derivabilidade and Consistency

We will now establish a number of properties of the **derivabilidade** relation. They are independently interesting, but each will play a role in the proof of the completeness theorem.

*fol:tab:prv:
sec*

Proposição tab.18. If $\Gamma \vdash \varphi$ and $\Gamma \cup \{\varphi\}$ is inconsistent, then Γ is inconsistent.

*fol:tab:prv:
prop:provability-contr*

Prova. There are finite $\Gamma_0 = \{\psi_1, \dots, \psi_n\}$ and $\Gamma_1 = \{\chi_1, \dots, \chi_n\} \subseteq \Gamma$ such that

$$\begin{aligned} &\{\mathbb{F} \varphi, \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\} \\ &\{\mathbb{T} \varphi, \mathbb{T} \chi_1, \dots, \mathbb{T} \chi_m\} \end{aligned}$$

have closed **tableaux**. Using the Cut rule on φ we can combine these into a single closed **tableau** that shows $\Gamma_0 \cup \Gamma_1$ is inconsistent. Since $\Gamma_0 \subseteq \Gamma$ and $\Gamma_1 \subseteq \Gamma$, $\Gamma_0 \cup \Gamma_1 \subseteq \Gamma$, hence Γ is inconsistent. $\square$

fol:tab:prv:
prop:prov-inc

Proposição tab.19. $\Gamma \vdash \varphi$ iff $\Gamma \cup \{\neg\varphi\}$ is inconsistent.

Prova. First suppose $\Gamma \vdash \varphi$, O.s., there is a closed **tableau** for

$$\{\mathbb{F} \varphi, \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}$$

Using the $\neg\mathbb{T}$ rule, this can be turned into a closed **tableau** for

$$\{\mathbb{T} \neg\varphi, \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}.$$

On the other hand, if there is a closed **tableau** for the latter, we can turn it into a closed **tableau** of the former by removing every formula that results from $\neg\mathbb{T}$ applied to the first assumption $\mathbb{T} \neg\varphi$ as well as that assumption, and adding the assumption $\mathbb{F} \varphi$. For if a branch was closed before because it contained the conclusion of $\neg\mathbb{T}$ applied to $\mathbb{T} \neg\varphi$, O.s., $\mathbb{F} \varphi$, the corresponding branch in the new **tableau** is also closed. If a branch in the old tableau was closed because it contained the assumption $\mathbb{T} \neg\varphi$ as well as $\mathbb{F} \neg\varphi$ we can turn it into a closed branch by applying $\neg\mathbb{F}$ to $\mathbb{F} \neg\varphi$ to obtain $\mathbb{T} \varphi$. This closes the branch since we added $\mathbb{F} \varphi$ as an assumption. $\square$

Problema tab.7. Prove that $\Gamma \vdash \neg\varphi$ iff $\Gamma \cup \{\varphi\}$ is inconsistent.

fol:tab:prv:
prop:explicit-inc

Proposição tab.20. If $\Gamma \vdash \varphi$ and $\neg\varphi \in \Gamma$, then Γ is inconsistent.

Prova. Suppose $\Gamma \vdash \varphi$ and $\neg\varphi \in \Gamma$. Then there are $\psi_1, \dots, \psi_n \in \Gamma$ such that

$$\{\mathbb{F} \varphi, \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}$$

has a closed tableau. Replace the assumption $\mathbb{F} \varphi$ by $\mathbb{T} \neg\varphi$, and insert the conclusion of $\neg\mathbb{T}$ applied to $\mathbb{F} \varphi$ after the assumptions. Any **sentença** in the **tableau** justified by appeal to line 1 in the old **tableau** is now justified by appeal to line $n+1$. So if the old **tableau** was closed, the new one is. It shows that Γ is inconsistent, since all assumptions are in Γ . $\square$

fol:tab:prv:
prop:provability-exhaustive

Proposição tab.21. If $\Gamma \cup \{\varphi\}$ and $\Gamma \cup \{\neg\varphi\}$ are both inconsistent, then Γ is inconsistent.

Prova. If there are $\psi_1, \dots, \psi_n \in \Gamma$ and $\chi_1, \dots, \chi_m \in \Gamma$ such that

$$\{\mathbb{T}\varphi, \mathbb{T}\psi_1, \dots, \mathbb{T}\psi_n\} \text{ and } \{\mathbb{T}\neg\varphi, \mathbb{T}\chi_1, \dots, \mathbb{T}\chi_m\}$$

both have closed **tableaux**, we can construct a single, combined **tableau** that shows that Γ is inconsistent by using as assumptions $\mathbb{T}\psi_1, \dots, \mathbb{T}\psi_n$ together with $\mathbb{T}\chi_1, \dots, \mathbb{T}\chi_m$, followed by an application of the Cut rule. This yields two branches, one starting with $\mathbb{T}\varphi$, the other with $\mathbb{F}\varphi$.

On the left left side, add the part of the first **tableau** below its assumptions. Here, every rule application is still correct, since each of the assumptions of the first **tableau**, including $\mathbb{T}\varphi$, is available. Thus, every branch below $\mathbb{T}\varphi$ closes.

On the right side, add the part of the second **tableau** below its assumption, with the results of any applications of $\neg\mathbb{T}$ to $\mathbb{T}\neg\varphi$ removed. The conclusion of $\neg\mathbb{T}$ to $\mathbb{T}\neg\varphi$ is $\mathbb{F}\varphi$, which is nevertheless available, as it is the conclusion of the Cut rule on the right side of the combined **tableau**.

If a branch in the second tableau was closed because it contained the assumption $\mathbb{T}\neg\varphi$ (which no longer appears as an assumption in the combined **tableau**) as well as $\mathbb{F}\neg\varphi$, we can applying $\neg\mathbb{F}$ to $\mathbb{F}\neg\varphi$ to obtain $\mathbb{T}\varphi$. Now the corresponding branch in the combined **tableau** also closes, because it contains the right-hand conclusion of the Cut rule, $\mathbb{F}\varphi$. If a branch in the second **tableau** closed for any other reason, the corresponding branch in the combined **tableau** also closes, since any **fórmulas assinadas** other than $\mathbb{T}\neg\varphi$ occurring on the branch in the old, second **tableau** also occur on the corresponding branch in the combined **tableau**. $\square$

-NoValue-

tab.9 Derivabilidade and the Propositional Connectives

explanation

We establish that the **derivabilidade** relation $\vdash$ of tableaux is strong enough to establish some basic facts involving the propositional connectives, such as that $\varphi \wedge \psi \vdash \varphi$ and $\varphi, \varphi \rightarrow \psi \vdash \psi$ (modus ponens). These facts are needed for the proof of the completeness theorem.

fol:tab:prr:
sec

Proposição tab.22.

1. Both $\varphi \wedge \psi \vdash \varphi$ and $\varphi \wedge \psi \vdash \psi$.
2. $\varphi, \psi \vdash \varphi \wedge \psi$.

fol:tab:prr:
prop:provability-land
fol:tab:prr:
prop:provability-land-left
fol:tab:prr:
prop:provability-land-right

Prova. 1. Both $\{\mathbb{F}\varphi, \mathbb{T}\varphi \wedge \psi\}$ and $\{\mathbb{F}\psi, \mathbb{T}\varphi \wedge \psi\}$ have closed **tableaux**

1.	$\mathbb{F}\varphi$	Assumption
2.	$\mathbb{T}\varphi \wedge \psi$	Assumption
3.	$\mathbb{T}\varphi$	$\wedge\mathbb{T}2$
4.	$\mathbb{T}\psi$	$\wedge\mathbb{T}2$
	$\otimes$	

- | | | |
|----|---------------------------------|----------------------|
| 1. | $\mathbb{F}\psi$ | Assumption |
| 2. | $\mathbb{T}\varphi \wedge \psi$ | Assumption |
| 3. | $\mathbb{T}\varphi$ | $\wedge\mathbb{T} 2$ |
| 4. | $\mathbb{T}\psi$ | $\wedge\mathbb{T} 2$ |
| | $\otimes$ | |

2. Here is a closed **tableau** for $\{\mathbb{T}\varphi, \mathbb{T}\psi, \mathbb{F}\varphi \wedge \psi\}$:

- | | | |
|----|--|----------------------|
| 1. | $\mathbb{F}\varphi \wedge \psi$ | Assumption |
| 2. | $\mathbb{T}\varphi$ | Assumption |
| 3. | $\mathbb{T}\psi$ | Assumption |
| | $\swarrow \quad \searrow$ | |
| 4. | $\mathbb{F}\varphi \quad \mathbb{F}\psi$ | $\wedge\mathbb{F} 1$ |
| | $\otimes \quad \otimes$ | |

fol:tab:pvr: **Proposição tab.23.**
prop:provability-lor

1. $\{\varphi \vee \psi, \neg\varphi, \neg\psi\}$ is inconsistent.
2. Both $\varphi \vdash \varphi \vee \psi$ and $\psi \vdash \varphi \vee \psi$.

Prova. 1. We give a closed **tableau** of $\{\mathbb{T}\varphi \vee \psi, \mathbb{T}\neg\varphi, \mathbb{T}\neg\psi\}$:

- | | | |
|----|--|--------------------|
| 1. | $\mathbb{T}\varphi \vee \psi$ | Assumption |
| 2. | $\mathbb{T}\neg\varphi$ | Assumption |
| 3. | $\mathbb{T}\neg\psi$ | Assumption |
| 4. | $\mathbb{F}\varphi$ | $\neg\mathbb{T} 2$ |
| 5. | $\mathbb{F}\psi$ | $\neg\mathbb{T} 3$ |
| | $\swarrow \quad \searrow$ | |
| 6. | $\mathbb{T}\varphi \quad \mathbb{T}\psi$ | $\vee\mathbb{T} 1$ |
| | $\otimes \quad \otimes$ | |

2. Both $\{\mathbb{F}\varphi \vee \psi, \mathbb{T}\varphi\}$ and $\{\mathbb{F}\varphi \vee \psi, \mathbb{T}\psi\}$ have closed **tableaux**:

- | | | |
|----|-------------------------------|--------------------|
| 1. | $\mathbb{F}\varphi \vee \psi$ | Assumption |
| 2. | $\mathbb{T}\varphi$ | Assumption |
| 3. | $\mathbb{F}\varphi$ | $\vee\mathbb{F} 1$ |
| 4. | $\mathbb{F}\psi$ | $\vee\mathbb{F} 1$ |
| | $\otimes$ | |

1.	$\mathbb{F} \varphi \vee \psi$	Assumption
2.	$\mathbb{T} \psi$	Assumption
3.	$\mathbb{F} \varphi$	$\vee \mathbb{F} 1$
4.	$\mathbb{F} \psi$	$\vee \mathbb{F} 1$
	$\otimes$	

Proposição tab.24.

*fol:tab:ppr:
prop:provability-lif*

1. $\varphi, \varphi \rightarrow \psi \vdash \psi$.

*fol:tab:ppr:
prop:provability-lif-left*

2. Both $\neg\varphi \vdash \varphi \rightarrow \psi$ and $\psi \vdash \varphi \rightarrow \psi$.

*fol:tab:ppr:
prop:provability-lif-right*

Prova. 1. $\{\mathbb{F} \psi, \mathbb{T} \varphi \rightarrow \psi, \mathbb{T} \varphi\}$ has a closed **tableau**:

1.	$\mathbb{F} \psi$	Assumption
2.	$\mathbb{T} \varphi \rightarrow \psi$	Assumption
3.	$\mathbb{T} \varphi$	Assumption
	$\swarrow \quad \searrow$	
4.	$\mathbb{F} \varphi \quad \mathbb{T} \psi$	$\rightarrow \mathbb{T} 2$
	$\otimes \quad \otimes$	

2. Both $\{\mathbb{F} \varphi \rightarrow \psi, \mathbb{T} \neg\varphi\}$ and $\{\mathbb{F} \varphi \rightarrow \psi, \mathbb{T} \psi\}$ have closed **tableaux**:

1.	$\mathbb{F} \varphi \rightarrow \psi$	Assumption
2.	$\mathbb{T} \neg\varphi$	Assumption
3.	$\mathbb{T} \varphi$	$\rightarrow \mathbb{F} 1$
4.	$\mathbb{F} \psi$	$\rightarrow \mathbb{F} 1$
5.	$\mathbb{F} \varphi$	$\neg \mathbb{T} 2$
	$\otimes$	

1.	$\mathbb{F} \varphi \rightarrow \psi$	Assumption
2.	$\mathbb{T} \psi$	Assumption
3.	$\mathbb{T} \varphi$	$\rightarrow \mathbb{F} 1$
4.	$\mathbb{F} \psi$	$\rightarrow \mathbb{F} 1$
	$\otimes$	

-NoValue-

tab.10 Derivabilidade and the Quantifiers

fol:tab:qpr: sec The completeness theorem also requires that the tableaux rules yield the facts explanation about $\vdash$ established in this section.

fol:tab:qpr: thm:strong-generalization **Teorema tab.25.** *If c is a constant not occurring in Γ or $\varphi(x)$ and $\Gamma \vdash \varphi(c)$, then $\Gamma \vdash \forall x \varphi(x)$.*


Prova. Suppose $\Gamma \vdash \varphi(c)$, O.s., there are $\psi_1, \dots, \psi_n \in \Gamma$ and a closed **tableau** for

$$\{\mathbb{F} \varphi(c), \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}.$$

We have to show that there is also a closed **tableau** for

$$\{\mathbb{F} \forall x \varphi(x), \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}.$$

Take the closed **tableau** and replace the first assumption with $\mathbb{F} \forall x \varphi(x)$, and insert $\mathbb{F} \varphi(c)$ after the assumptions.



The tableau is still closed, since all **sentenças** available as assumptions before are still available at the top of the **tableau**. The inserted line is the result of a correct application of $\forall\mathbb{F}$, since the **símbolo de constante** c does not occur in $\psi_1, \dots, \psi_n$ or $\forall x \varphi(x)$, O.s., it does not occur above the inserted line in the new **tableau**. $\square$

fol:tab:qpr: prop:provability-quantifiers **Proposição tab.26.**

1. $\varphi(t) \vdash \exists x \varphi(x)$.
2. $\forall x \varphi(x) \vdash \varphi(t)$.

Prova. 1. A closed **tableau** for $\mathbb{F} \exists x \varphi(x), \mathbb{T} \varphi(t)$ is:

- | | | |
|----|-----------------------------------|-----------------------|
| 1. | $\mathbb{F} \exists x \varphi(x)$ | Assumption |
| 2. | $\mathbb{T} \varphi(t)$ | Assumption |
| 3. | $\mathbb{F} \varphi(t)$ | $\exists\mathbb{F} 1$ |
| | $\otimes$ | |

2. A closed **tableau** for $\mathbb{F} \varphi(t), \mathbb{T} \forall x \varphi(x)$, is:

- | | | |
|----|-----------------------------------|------------------------|
| 1. | $\mathbb{F} \varphi(t)$ | Assumption |
| 2. | $\mathbb{T} \forall x \varphi(x)$ | Assumption |
| 3. | $\mathbb{T} \varphi(t)$ | $\forall \mathbb{T} 2$ |
| | $\otimes$ | |

-NoValue-

tab.11 Soundness

explanation A *derivação* system, such as tableaux, is *sound* if it cannot *deriva* things that do not actually hold. Soundness is thus a kind of guaranteed safety property for *derivação* systems. Depending on which proof theoretic property is in question, we would like to know for instance, that fol:tab:sou:sec

1. every *derivável* φ is valid;
2. if a *sentença* is *derivável* from some others, it is also a consequence of them;
3. if a set of *sentenças* is inconsistent, it is unsatisfiable.

These are important properties of a *derivação* system. If any of them do not hold, the *derivação* system is deficient—it would *deriva* too much. Consequently, establishing the soundness of a *derivação* system is of the utmost importance.

Because all these proof-theoretic properties are defined via closed *tableaux* of some kind or other, proving (1)–(3) above requires proving something about the semantic properties of closed *tableaux*. We will first define what it means for a *fórmula assinada* to be satisfied in a structure, and then show that if a *tableau* is closed, no structure satisfies all its assumptions. (1)–(3) then follow as corollaries from this result.

Definição tab.27. A *estrutura* $\mathfrak{M}$ *satisfies* a *fórmula assinada* $\mathbb{T} \varphi$ iff $\mathfrak{M} \models \varphi$, and it satisfies $\mathbb{F} \varphi$ iff $\mathfrak{M} \not\models \varphi$. $\mathfrak{M}$ satisfies a set of *fórmulas assinadas* Γ iff it satisfies every $S \varphi \in \Gamma$. Γ is *satisfiable* if there is a *estrutura* that satisfies it, and *unsatisfiable* otherwise.

Teorema tab.28 (Soundness). *If Γ has a closed *tableau*, Γ is unsatisfiable.* fol:tab:sou:thm:tableau-soundness

Prova. Let's call a branch of a *tableau* satisfiable iff the set of *fórmulas assinadas* on it is satisfiable, and let's call a *tableau* satisfiable if it contains at least one satisfiable branch.

We show the following: Extending a satisfiable *tableau* by one of the rules of inference always results in a satisfiable *tableau*. This will prove the theorem: any closed *tableau* results by applying rules of inference to the *tableau* consisting only of assumptions from Γ . So if Γ were satisfiable, any *tableau* for it would be satisfiable. A closed *tableau*, however, is clearly not satisfiable:

every branch contains both $\mathbb{T}\varphi$ and $\mathbb{F}\varphi$, and no structure can both satisfy and not satisfy φ .

Suppose we have a satisfiable **tableau**, O.s., a **tableau** with at least one satisfiable branch. Applying a rule of inference either adds **fórmulas assinadas** to a branch, or splits a branch in two. If the **tableau** has a satisfiable branch which is not extended by the rule application in question, it remains a satisfiable branch in the extended **tableau**, so the extended tableau is satisfiable. So we only have to consider the case where a rule is applied to a satisfiable branch.

Let Γ be the set of **fórmulas assinadas** on that branch, and let $S\varphi \in \Gamma$ be the **fórmula assinada** to which the rule is applied. If the rule does not result in a split branch, we have to show that the extended branch, O.s., Γ together with the conclusions of the rule, is still satisfiable. If the rule results in a split branch, we have to show that at least one of the two resulting branches is satisfiable.

First, we consider the possible inferences that do not result in a split branch.

1. The branch is expanded by applying $\neg\mathbb{T}$ to $\mathbb{T}\neg\psi \in \Gamma$. Then the extended branch contains the **fórmulas assinadas** $\Gamma \cup \{\mathbb{F}\psi\}$. Suppose $\mathfrak{M} \models \Gamma$. In particular, $\mathfrak{M} \models \neg\psi$. Thus, $\mathfrak{M} \not\models \psi$, O.s., $\mathfrak{M}$ satisfies $\mathbb{F}\psi$.
2. The branch is expanded by applying $\neg\mathbb{F}$ to $\mathbb{F}\neg\psi \in \Gamma$: Exercise.
3. The branch is expanded by applying $\wedge\mathbb{T}$ to $\mathbb{T}\psi \wedge \chi \in \Gamma$, which results in two new **fórmulas assinadas** on the branch: $\mathbb{T}\psi$ and $\mathbb{T}\chi$. Suppose $\mathfrak{M} \models \Gamma$, in particular $\mathfrak{M} \models \psi \wedge \chi$. Then $\mathfrak{M} \models \psi$ and $\mathfrak{M} \models \chi$. This means that $\mathfrak{M}$ satisfies both $\mathbb{T}\psi$ and $\mathbb{T}\chi$.
4. The branch is expanded by applying $\vee\mathbb{F}$ to $\mathbb{F}\psi \vee \chi \in \Gamma$: Exercise.
5. The branch is expanded by applying $\rightarrow\mathbb{F}$ to $\mathbb{F}\psi \rightarrow \chi \in \Gamma$: This results in two new **fórmulas assinadas** on the branch: $\mathbb{T}\psi$ and $\mathbb{F}\chi$. Suppose $\mathfrak{M} \models \Gamma$, in particular $\mathfrak{M} \models \psi \rightarrow \chi$. Then $\mathfrak{M} \models \psi$ and $\mathfrak{M} \models \chi$. This means that $\mathfrak{M}$ satisfies both $\mathbb{T}\psi$ and $\mathbb{F}\chi$.
6. The branch is expanded by applying $\forall\mathbb{T}$ to $\mathbb{T}\forall x\psi(x) \in \Gamma$: This results in a new **fórmula assinada** $\mathbb{T}\varphi(t)$ on the branch. Suppose $\mathfrak{M} \models \Gamma$, in particular, $\mathfrak{M} \models \forall x\psi(x)$. By ??, $\mathfrak{M} \models \varphi(t)$. Consequently, $\mathfrak{M}$ satisfies $\mathbb{T}\varphi(t)$.
7. The branch is expanded by applying $\forall\mathbb{F}$ to $\mathbb{F}\forall x\psi(x) \in \Gamma$: This results in a new **fórmula assinada** $\mathbb{F}\varphi(a)$ where a is a **símbolo de constante** not occurring in Γ . Since Γ is satisfiable, there is a $\mathfrak{M}$ such that $\mathfrak{M} \models \Gamma$, in particular $\mathfrak{M} \models \forall x\psi(x)$. We have to show that $\Gamma \cup \{\mathbb{F}\varphi(a)\}$ is satisfiable. To do this, we define a suitable $\mathfrak{M}'$ as follows.

By ??, $\mathfrak{M} \models \forall x\psi(x)$ iff for some s , $\mathfrak{M}, s \models \psi(x)$. Now let $\mathfrak{M}'$ be just like $\mathfrak{M}$, except $a^{\mathfrak{M}'} = s(x)$. By ??, for any $\mathbb{T}\chi \in \Gamma$, $\mathfrak{M}' \models \chi$, and for any $\mathbb{F}\chi \in \Gamma$, $\mathfrak{M}' \not\models \chi$, since a does not occur in Γ .

By ??, $\mathfrak{M}', s \not\models \varphi(x)$. By ??, $\mathfrak{M}', s \not\models \varphi(a)$. Since $\varphi(a)$ is a **sentença**, by ??, $\mathfrak{M}' \not\models \varphi(a)$, O.s., $\mathfrak{M}'$ satisfies $\mathbb{F} \varphi(a)$.

8. The branch is expanded by applying $\exists\mathbb{T}$ to $\mathbb{T} \exists x \psi(x) \in \Gamma$: Exercise.

9. The branch is expanded by applying $\exists\mathbb{F}$ to $\mathbb{F} \exists x \psi(x) \in \Gamma$: Exercise.

Now let's consider the possible inferences that result in a split branch.

1. The branch is expanded by applying $\wedge\mathbb{F}$ to $\mathbb{F} \psi \wedge \chi \in \Gamma$, which results in two branches, a left one continuing through $\mathbb{F} \psi$ and a right one through $\mathbb{F} \chi$. Suppose $\mathfrak{M} \models \Gamma$, in particular $\mathfrak{M} \not\models \psi \wedge \chi$. Then $\mathfrak{M} \not\models \psi$ or $\mathfrak{M} \not\models \chi$. In the former case, $\mathfrak{M}$ satisfies $\mathbb{F} \psi$, O.s., $\mathfrak{M}$ satisfies the formulas on the left branch. In the latter, $\mathfrak{M}$ satisfies $\mathbb{F} \chi$, O.s., $\mathfrak{M}$ satisfies the formulas on the right branch.
2. The branch is expanded by applying $\vee\mathbb{T}$ to $\mathbb{T} \psi \vee \chi \in \Gamma$: Exercise.
3. The branch is expanded by applying $\rightarrow\mathbb{T}$ to $\mathbb{T} \psi \rightarrow \chi \in \Gamma$: Exercise.
4. The branch is expanded by Cut: This results in two branches, one containing $\mathbb{T} \psi$, the other containing $\mathbb{F} \psi$. Since $\mathfrak{M} \models \Gamma$ and either $\mathfrak{M} \models \psi$ or $\mathfrak{M} \not\models \psi$, $\mathfrak{M}$ satisfies either the left or the right branch. $\square$

Problema tab.8. Complete the proof of **Teorema tab.28**.

Corolário tab.29. *If $\vdash \varphi$ then φ is valid.*

*fol:tab:sou:
cor:weak-soundness*

Corolário tab.30. *If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$.*

*fol:tab:sou:
cor:entailment-soundness*

Prova. If $\Gamma \vdash \varphi$ then for some $\psi_1, \dots, \psi_n \in \Gamma$, $\{\mathbb{F} \varphi, \mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}$ has a closed **tableau**. By **Teorema tab.28**, every **estrutura** $\mathfrak{M}$ either makes some ψ_i false or makes φ true. Hence, if $\mathfrak{M} \models \Gamma$ then also $\mathfrak{M} \models \varphi$. $\square$

Corolário tab.31. *If Γ is satisfiable, then it is consistent.*

*fol:tab:sou:
cor:consistency-soundness*

Prova. We prove the contrapositive. Suppose that Γ is not consistent. Then there are $\psi_1, \dots, \psi_n \in \Gamma$ and a closed **tableau** for $\{\mathbb{T} \psi_1, \dots, \mathbb{T} \psi_n\}$. By **Teorema tab.28**, there is no $\mathfrak{M}$ such that $\mathfrak{M} \models \psi_i$ for all $i = 1, \dots, n$. But then Γ is not satisfiable. $\square$

-NoValue-

tab.12 Tableaux with Predicado de identidade

Tableaux with **predicado de identidade** require additional inference rules. The rules for $=$ are (t, t_1 , and t_2 are closed terms):

*fol:tab:ide:
sec*

$\frac{}{\mathbb{T} t = t} =$	$\frac{\mathbb{T} t_1 = t_2 \quad \mathbb{T} \varphi(t_1)}{\mathbb{T} \varphi(t_2)} = \mathbb{T}$	$\frac{\mathbb{T} t_1 = t_2 \quad \mathbb{F} \varphi(t_1)}{\mathbb{F} \varphi(t_2)} = \mathbb{F}$
-------------------------------	---	---

Note that in contrast to all the other rules, $=\mathbb{T}$ and $=\mathbb{F}$ require that *two* signed **fórmulas** already appear on the branch, namely both $\mathbb{T} t_1 = t_2$ and $\mathbb{F} \varphi(t_1)$.

Exemplo tab.32. If s and t are closed terms, then $s = t, \varphi(s) \vdash \varphi(t)$:

1. $\mathbb{F} \varphi(t)$ Assumption
 2. $\mathbb{T} s = t$ Assumption
 3. $\mathbb{T} \varphi(s)$ Assumption
 4. $\mathbb{T} \varphi(t)$ $=\mathbb{T} 2, 3$
- $\otimes$

This may be familiar as the principle of substitutability of identicals, or Leibniz' Law.

Tableaux prove that $=$ is symmetric, O.s., that $s_1 = s_2 \vdash s_2 = s_1$:

1. $\mathbb{F} s_2 = s_1$ Assumption
 2. $\mathbb{T} s_1 = s_2$ Assumption
 3. $\mathbb{T} s_1 = s_1$ $=$
 4. $\mathbb{T} s_2 = s_1$ $=\mathbb{T} 2, 3$
- $\otimes$

Here, line 2 is the first prerequisite **fórmula** $\mathbb{T} s_1 = s_2$ of $=\mathbb{T}$. Line 3 is the second one, of the form $\mathbb{T} \varphi(s_2)$ —think of $\varphi(x)$ as $x = s_1$, then $\varphi(s_1)$ is $s_1 = s_1$ and $\varphi(s_2)$ is $s_2 = s_1$.

They also prove that $=$ is transitive, O.s., that $s_1 = s_2, s_2 = s_3 \vdash s_1 = s_3$:

1. $\mathbb{F} s_1 = s_3$ Assumption
 2. $\mathbb{T} s_1 = s_2$ Assumption
 3. $\mathbb{T} s_2 = s_3$ Assumption
 4. $\mathbb{T} s_1 = s_3$ $=\mathbb{T} 3, 2$
- $\otimes$

In this **tableau**, the first prerequisite **fórmula** of $=\mathbb{T}$ is line 3, $\mathbb{T} s_2 = s_3$ (s_2 plays the role of t_1 , and s_3 the role of t_2). The second prerequisite, of the form $\mathbb{T} \varphi(s_2)$ is line 2. Here, think of $\varphi(x)$ as $s_1 = x$; that makes $\varphi(s_2)$ into $t_1 = t_2$ (O.s., line 2) and $\varphi(s_3)$ into the **fórmula** $s_1 = s_3$ in the conclusion.

Problema tab.9. Give closed **tableaux** for the following:

1. $\mathbb{F} \forall x \forall y ((x = y \wedge \varphi(x)) \rightarrow \varphi(y))$

2. $\mathbb{F} \exists x (\varphi(x) \wedge \forall y (\varphi(y) \rightarrow y = x)),$
 $\mathbb{T} \exists x \varphi(x) \wedge \forall y \forall z ((\varphi(y) \wedge \varphi(z)) \rightarrow y = z)$

-NoValue-

tab.13 Soundness with Predicado de identidade

Proposição tab.33. *Tableaux with rules for identity are sound: no closed tableau is satisfiable.*

fol:tab:sid:
sec

Prova. We just have to show as before that if a tableau has a satisfiable branch, the branch resulting from applying one of the rules for = to it is also satisfiable. Let Γ be the set of signed fórmulas on the branch, and let $\mathfrak{M}$ be a estrutura satisfying Γ .

Suppose the branch is expanded using =, O.s., by adding the signed fórmula $\mathbb{T} t = t$. Trivially, $\mathfrak{M} \models t = t$, so $\mathfrak{M}$ also satisfies $\Gamma \cup \{\mathbb{T} t = t\}$.

If the branch is expanded using $=\mathbb{T}$, we add a signed fórmula $S \varphi(t_2)$, but Γ contains both $\mathbb{T} t_1 = t_2$ and $\mathbb{T} \varphi(t_1)$. Thus we have $\mathfrak{M} \models t_1 = t_2$ and $\mathfrak{M} \models \varphi(t_1)$. Let s be a variable assignment with $s(x) = \text{Val}^{\mathfrak{M}}(t_1)$. By ??, $\mathfrak{M}, s \models \varphi(t_1)$. Since $s \sim_x s$, by ??, $\mathfrak{M}, s \models \varphi(x)$. since $\mathfrak{M} \models t_1 = t_2$, we have $\text{Val}^{\mathfrak{M}}(t_1) = \text{Val}^{\mathfrak{M}}(t_2)$, and hence $s(x) = \text{Val}^{\mathfrak{M}}(t_2)$. By applying ?? again, we also have $\mathfrak{M}, s \models \varphi(t_2)$. By ??, $\mathfrak{M} \models \varphi(t_2)$. The case of $=\mathbb{F}$ is treated similarly. $\square$

Créditos das Imagens

Referências Bibliográficas