

-NoValue-

cr.1 β -reduction

lam:cr:b:
sec
lam:cr:b:
lem:one-par

Lema cr.1. If $M \xrightarrow{\beta} M'$, then $M \xRightarrow{\beta} M'$.

Prova. If $M \xrightarrow{\beta} M'$, then M is $(\lambda x. N)Q$, M' is $N[Q/x]$, for some x , N , and Q . Since $N \xRightarrow{\beta} N$ and $Q \xRightarrow{\beta} Q$ by ??, we immediately have $(\lambda x. N)Q \xRightarrow{\beta} N[Q/x]$ by ????. $\square$

lam:cr:b:
lem:par-red

Lema cr.2. If $M \xRightarrow{\beta} M'$, then $M \xrightarrow{\beta} M'$.

Prova. By induction on the *derivação* of $M \xRightarrow{\beta} M'$.

1. The last rule is ?? : Then M and M' are just x , and $x \xrightarrow{\beta} x$.
2. The last rule is ?? : M is $\lambda x. N$ and M' is $\lambda x. N'$ for some x , N , N' , where $N \xRightarrow{\beta} N'$. By induction hypothesis we have $N \xrightarrow{\beta} N'$. Then $\lambda x. N \xrightarrow{\beta} \lambda x. N'$ (by the same series of $\xrightarrow{\beta}$ contractions as $N \xrightarrow{\beta} N'$).
3. The last rule is ?? : M is PQ and M' is $P'Q'$ for some P , Q , P' , Q' , where $P \xRightarrow{\beta} P'$ and $Q \xRightarrow{\beta} Q'$. By induction hypothesis we have $P \xrightarrow{\beta} P'$ and $Q \xrightarrow{\beta} Q'$. So $PQ \xrightarrow{\beta} P'Q'$ by the reduction sequence $P \xrightarrow{\beta} P'$ followed by the reduction $Q \xrightarrow{\beta} Q'$.
4. The last rule is ?? : M is $(\lambda x. N)Q$ and M' is $N'[Q'/x]$ for some x , N , M' , Q , Q' , where $N \xRightarrow{\beta} N'$ and $Q \xRightarrow{\beta} Q'$. By induction hypothesis we get $Q \xrightarrow{\beta} Q'$ and $N \xrightarrow{\beta} N'$. So $(\lambda x. N)Q \xrightarrow{\beta} N'[Q'/x]$ by $N \xrightarrow{\beta} N'$ followed by $Q \xrightarrow{\beta} Q'$ and finally contraction of $(\lambda x. N')Q'$ to $N'[Q'/x]$. $\square$

lam:cr:b:
lem:str

Lema cr.3. $\xrightarrow{\beta}$ is the smallest transitive relation containing $\xRightarrow{\beta}$.

Prova. Let $\xrightarrow{X}$ be the smallest transitive relation containing $\xRightarrow{\beta}$.

$\xrightarrow{\beta} \subseteq \xrightarrow{X}$: Suppose $M \xrightarrow{\beta} M'$, O.s., $M \equiv M_1 \xrightarrow{\beta} \dots \xrightarrow{\beta} M_k \equiv M'$. By **Lema cr.1**, $M \equiv M_1 \xRightarrow{\beta} \dots \xRightarrow{\beta} M_k \equiv M'$. Since $\xrightarrow{X}$ contains $\xRightarrow{\beta}$ and is transitive, $M \xrightarrow{X} M'$.

$\xrightarrow{X} \subseteq \xrightarrow{\beta}$: Suppose $M \xrightarrow{X} M'$, O.s., $M \equiv M_1 \xRightarrow{\beta} \dots \xRightarrow{\beta} M_k \equiv M'$. By **Lema cr.2**, $M \equiv M_1 \xrightarrow{\beta} \dots \xrightarrow{\beta} M_k \equiv M'$. Since $\xrightarrow{\beta}$ is transitive, $M \xrightarrow{\beta} M'$. $\square$

lam:cr:b:
thm:cr

Teorema cr.4. $\xrightarrow{\beta}$ satisfies the Church–Rosser property.

Prova. Immediate from ??, ??, and **Lema cr.3**.

□

Créditos das Imagens

Referências Bibliográficas