

-NoValue-

## syn.1 The De Bruijn Index

lam:syn:deb:  
sec

$\alpha$ -Equivalence is very natural, as terms that are  $\alpha$ -equivalent “mean the same.” In fact, it is possible to give a syntax for lambda terms which does not distinguish terms that can be  $\alpha$ -converted to each other. The best known replaces variables by their *De Bruijn index*.

When we write  $\lambda x. M$ , we explicitly state that  $x$  is the parameter of the function, so that we can use  $x$  in  $M$  to refer to this parameter. In the de Bruijn index, however, parameters have no name and reference to them in the function body is denoted by a number denoting the levels of abstraction between them. For example, consider the example of  $\lambda x. \lambda y. yx$ : the outer abstraction is on binds the variable  $x$ ; the inner abstraction binds the variable is  $y$ ; the sub-term  $yx$  lies in the scope of the inner abstraction: there is no abstraction between  $y$  and its abstract  $\lambda y$ , but one abstract between  $x$  and its abstract  $\lambda x$ . Thus we write 0 1 for  $yx$ , and  $\lambda. \lambda. 01$  for the entire term.

**Definição syn.1.** De Bruijn terms are inductively defines as follows:

1.  $n$ , where  $n$  is any natural number.
2.  $PQ$ , where  $P$  and  $Q$  are both De Bruijn terms.
3.  $\lambda. N$ , where  $N$  is a De Bruijn term.

A formalized translation from ordinary lambda terms to De Bruijn indexed terms is as follows:

**Definição syn.2.**

$$\begin{aligned} F_\Gamma(x) &= \Gamma(x) \\ F_\Gamma(PQ) &= F_\Gamma(P)F_\Gamma(Q) \\ F_\Gamma(\lambda x. N) &= \lambda. F_{x,\Gamma}(N) \end{aligned}$$

where  $\Gamma$  is a list of variables indexed from zero, and  $\Gamma(x)$  denotes the position of the variable  $x$  in  $\Gamma$ . For example, if  $\Gamma$  is  $x, y, z$ , then  $\Gamma(x)$  is 0 and  $\Gamma(z)$  is 2.

$x, \Gamma$  denotes the list resulted from pushing  $x$  to the head of  $\Gamma$ ; for instance, continuing the  $\Gamma$  in last example,  $w, \Gamma$  is  $w, x, y, z$ .

Recovering a standard lambda term from a de Bruijn term is done as follows:

**Definição syn.3.**

$$\begin{aligned} G_\Gamma(n) &= \Gamma[n] \\ G_\Gamma(PQ) &= G_\Gamma(P)G_\Gamma(Q) \\ G_\Gamma(\lambda. N) &= \lambda x. G_{x,\Gamma}(N) \end{aligned}$$

where  $\Gamma$  is again a list of variables indexed from zero, and  $\Gamma[n]$  denotes the variable in position  $n$ . For example, if  $\Gamma$  is  $x, y, z$ , then  $\Gamma[1]$  is  $y$ .

The variable  $x$  in last equation is chosen to be any variable that not in  $\Gamma$ .

Here we give some results without proving them:

**Proposição syn.4.** *If  $M \xrightarrow{\alpha} M'$ , and  $\Gamma$  is any list containing  $\text{FV}(M)$ , then  $F_{\Gamma}(M) \equiv F_{\Gamma}(M')$ .*

## Créditos das Imagens

## Referências Bibliográficas