

-NoValue-

inf.1 Łukasiewicz logic

mv:inf:luk:
sec

This is a short “stub” of a section on infinite-valued Łukasiewicz logic.

mv:inf:luk:
def:lukasiewicz

Definição inf.1. Infinite-valued Łukasiewicz logic $\mathbf{L}_\infty$ is defined using the matrix:

1. The standard propositional language $\mathcal{L}_0$ with $\neg, \wedge, \vee, \rightarrow$.
2. The set of truth values V_∞ .
3. 1 is the only designated value, O.s., $V^+ = \{1\}$.
4. Truth functions are given by the following functions:

$$\begin{aligned}\neg_{\mathbf{L}}(x) &= 1 - x \\ \wedge_{\mathbf{L}}(x, y) &= \min(x, y) \\ \vee_{\mathbf{L}}(x, y) &= \max(x, y) \\ \rightarrow_{\mathbf{L}}(x, y) &= \min(1, 1 - (x - y)) = \begin{cases} 1 & \text{if } x \leq y \\ 1 - (x - y) & \text{otherwise.} \end{cases}\end{aligned}$$

m -valued Łukasiewicz logic is defined the same, except $V = V_m$.

Proposição inf.2. The logic $\mathbf{L}_3$ defined by ?? is the same as $\mathbf{L}_3$ defined by *Definição inf.1*.

Prova. This can be seen by comparing the truth tables for the connectives given in ?? with the truth tables determined by the equations in *Definição inf.1*:

$\neg$		$\wedge_{\mathbf{L}_3}$	1	1/2	0
1	0	1	1	1/2	0
1/2	1/2	1/2	1/2	1/2	0
0	1	0	0	0	0

$\vee_{\mathbf{L}_3}$	1	1/2	0	$\rightarrow_{\mathbf{L}_3}$	1	1/2	0
1	1	1	1	1	1	1/2	0
1/2	1	1/2	1/2	1/2	1	1	1/2
0	1	1/2	0	0	1	1	1

□

mv:inf:luk:
prop:luk-infty-m

Proposição inf.3. If $\Gamma \models_{\mathbf{L}_\infty} \psi$ then $\Gamma \models_{\mathbf{L}_m} \psi$ for all $m \geq 2$.

Prova. Exercise.

□

Problema inf.1. Prove **Proposição inf.3**.

In fact, the converse holds as well.

Infinite-valued Łukasiewicz logic is the most popular fuzzy logic. In the fuzzy logic literature, the conditional is often defined as $\neg\varphi \vee \psi$. The result would be an infinite-valued strong Kleene logic.

Problema inf.2. Show that $(p \rightarrow q) \vee (q \rightarrow p)$ is a tautology of $\mathbf{L}_\infty$.

Créditos das Imagens

Referências Bibliográficas