

-NoValue-

seq.1 Rules and Derivações

mv:seq:rul:
sec

For the following, let Γ, Δ, Π, A represent finite sequences of **sentenças**.

Definição seq.1 (Sequent). An *n-sided sequent* is an expression of the form

$$\Gamma_1 \mid \dots \mid \Gamma_n$$

where each Γ_i is a finite (possibly empty) sequences of **sentenças** of the language $\mathcal{L}$.

Definição seq.2 (Initial Sequent). An *n-sided initial sequent* is an *n-sided* sequent of the form $\varphi \mid \dots \mid \varphi$ for any **sentença** φ in the language.

If the language contains a 0-place connective $\star$, O.s., a propositional constant, then we also take the sequent $\dots \mid \star \mid \dots$ where $\star$ appears in the space for the truth value associated with $\tilde{\star} \in V$, and is empty otherwise.

For each connective of an *n-valued* logic $\mathbf{L}$, there is a logical rule for each truth value that this connective can take in $\mathbf{L}$. **Derivações** in an *n-sided* sequent calculus for $\mathbf{L}$ are trees of sequents, where the topmost sequents are initial sequents, and if a sequent stands below one or more other sequents, it must follow correctly by a rule of inference for the connectives of $\mathbf{L}$.

Definição seq.3 (Theorems). A **sentença** φ is a *theorem* of an *n-valued* logic $\mathbf{L}$ if there is a **derivação** of the *n*-sequent containing φ in each position corresponding to a designated truth value of $\mathbf{L}$. We write $\vdash_{\mathbf{L}} \varphi$ if φ is a theorem and $\not\vdash_{\mathbf{L}} \varphi$ if it is not.

Definição seq.4 (Derivabilidade). A **sentença** φ is *derivável* from a set of **sentenças** Γ in an *n-valued* logic $\mathbf{L}$, $\Gamma \vdash_{\mathbf{L}} \varphi$, iff there is a finite subset $\Gamma_0 \subseteq \Gamma$ and a sequence Γ'_0 of the **sentenças** in Γ_0 such that the following sequent has a **derivação**:

$$A_1 \mid \dots \mid A_n$$

where A_i is φ if position *i* corresponds to a designated truth value, and Γ'_0 otherwise. If φ is not *derivável* from Γ we write $\Gamma \not\vdash \varphi$.

For instance, 3-valued Łukasiewicz logic has a 3-sided sequent calculus. In a 3-sided sequent $\Gamma \mid \Pi \mid \Delta$, Γ corresponds to $\mathbb{F}$, Δ to $\mathbb{T}$, and Π to $\mathbb{U}$. Axioms are $\varphi \mid \varphi \mid \varphi$. Since only $\mathbb{T}$ is designated, $\Gamma \vdash_{\mathbf{L}_3} \varphi$ iff the sequent $\Gamma \mid \Gamma \mid \varphi$ has a **derivação**. (If $\mathbb{U}$ were also designated, we would need a **derivação** of $\Gamma \mid \varphi \mid \varphi$.)

Créditos das Imagens

Referências Bibliográficas