

-NoValue-

mar.1 Introduction

The *standard model* of arithmetic is the **estrutura** $\mathfrak{N}$ with $|\mathfrak{N}| = \mathbb{N}$ in which $\mathfrak{o}$, $\mathfrak{!}$, $+$, $\times$, and $<$ are interpreted as you would expect. That is, $\mathfrak{o}$ is 0, $\mathfrak{!}$ is the successor function, $+$ is interpreted as addition and $\times$ as multiplication of the numbers in $\mathbb{N}$. Specifically,

$$\begin{aligned}\mathfrak{o}^{\mathfrak{N}} &= 0 \\ \mathfrak{!}^{\mathfrak{N}}(n) &= n + 1 \\ +^{\mathfrak{N}}(n, m) &= n + m \\ \times^{\mathfrak{N}}(n, m) &= nm\end{aligned}$$

Of course, there are structures for $\mathcal{L}_A$ that have domains other than $\mathbb{N}$. For instance, we can take $\mathfrak{M}$ with domain $|\mathfrak{M}| = \{a\}^*$ (the finite sequences of the single symbol a , O.s., $\emptyset$, a , aa , aaa , $\dots$), and interpretations

$$\begin{aligned}\mathfrak{o}^{\mathfrak{M}} &= \emptyset \\ \mathfrak{!}^{\mathfrak{M}}(s) &= s \frown a \\ +^{\mathfrak{M}}(n, m) &= a^{n+m} \\ \times^{\mathfrak{M}}(n, m) &= a^{nm}\end{aligned}$$

These two structures are “essentially the same” in the sense that the only difference is the **membros** of the **domínios** but not how the **membros** of the **domínios** are related among each other by the interpretation functions. We say that the two **estructuras** are *isomorphic*.

It is an easy consequence of the compactness theorem that any theory true in $\mathfrak{N}$ also has models that are not isomorphic to $\mathfrak{N}$. Such structures are called *non-standard*. The interesting thing about them is that while the **membros** of a standard model (O.s., $\mathfrak{N}$, but also all **estructuras** isomorphic to it) are exhausted by the values of the standard numerals $\bar{n}$, O.s.,

$$|\mathfrak{N}| = \{\text{Val}^{\mathfrak{N}}(\bar{n}) : n \in \mathbb{N}\}$$

that isn’t the case in non-standard models: if $\mathfrak{M}$ is non-standard, then there is at least one $x \in |\mathfrak{M}|$ such that $x \neq \text{Val}^{\mathfrak{M}}(\bar{n})$ for all n .

These non-standard elements are pretty neat: they are “infinite natural numbers.” But their existence also explains, in a sense, the incompleteness phenomena. Consider an example, p.ex., the consistency statement for Peano arithmetic, $\text{Con}_{\mathbf{PA}}$, O.s., $\neg \exists x \text{Prf}_{\mathbf{PA}}(x, \ulcorner \perp \urcorner)$. Since $\mathbf{PA}$ neither proves $\text{Con}_{\mathbf{PA}}$ nor $\neg \text{Con}_{\mathbf{PA}}$, either can be consistently added to $\mathbf{PA}$. Since $\mathbf{PA}$ is consistent, $\mathfrak{N} \models \text{Con}_{\mathbf{PA}}$, and consequently $\mathfrak{N} \not\models \neg \text{Con}_{\mathbf{PA}}$. So $\mathfrak{N}$ is *not* a model of $\mathbf{PA} \cup \{\neg \text{Con}_{\mathbf{PA}}\}$, and all its models must be nonstandard. Models of

$\mathbf{PA} \cup \{\neg \text{Con}_{\mathbf{PA}}\}$ must contain some **membro** that serves as the witness that makes $\exists x \text{Prf}_{\mathbf{PA}}(\ulcorner \perp \urcorner)$ true, O.s., a Gödel number of a **derivação** of a contradiction from $\mathbf{PA}$. Such **um membro** can't be standard—since $\mathbf{PA} \vdash \neg \text{Prf}_{\mathbf{PA}}(\overline{n}, \ulcorner \perp \urcorner)$ for every n .

Créditos das Imagens

Referências Bibliográficas