

-NoValue-

prf.1 Proofs in Modal Systems

nml:prf:prs:
sec We now come to proofs in systems of modal logic other than **K**.

nml:prf:prs:
prop:S5facts **Proposição prf.1.** *The following provability results obtain:*

1. **KT5** $\vdash$ B;
2. **KT5** $\vdash$ 4;
3. **KDB4** $\vdash$ T;
4. **KB4** $\vdash$ 5;
5. **KB5** $\vdash$ 4;
6. **KT** $\vdash$ D.

nml:prf:prs:
prop:S5facts-KT-D

Prova. We exhibit proofs for each.

1. **KT5** $\vdash$ B:

1. **KT5** $\vdash \Diamond\varphi \rightarrow \Box\Diamond\varphi$ 5
2. **KT5** $\vdash \varphi \rightarrow \Diamond\varphi$ $T_\Diamond$
3. **KT5** $\vdash \varphi \rightarrow \Box\Diamond\varphi$ PL, 2, 1.

2. **KT5** $\vdash$ 4:

1. **KT5** $\vdash \Diamond\Box\varphi \rightarrow \Box\Diamond\Box\varphi$ 5 with $\Box\varphi$ for p
2. **KT5** $\vdash \Box\varphi \rightarrow \Diamond\Box\varphi$ $T_\Diamond$ with $\Box\varphi$ for p
3. **KT5** $\vdash \Box\varphi \rightarrow \Box\Diamond\Box\varphi$ PL, 2, 1
4. **KT5** $\vdash \Diamond\Box\varphi \rightarrow \Box\varphi$ $5_\Diamond$
5. **KT5** $\vdash \Box\Diamond\Box\varphi \rightarrow \Box\Box\varphi$ RK, 4
6. **KT5** $\vdash \Box\varphi \rightarrow \Box\Box\varphi$ PL, 3, 5.

3. **KDB4** $\vdash$ T:

1. **KDB4** $\vdash \Diamond\Box\varphi \rightarrow \varphi$ $B_\Diamond$
2. **KDB4** $\vdash \Box\Box\varphi \rightarrow \Diamond\Box\varphi$ D with $\Box\varphi$ for p
3. **KDB4** $\vdash \Box\Box\varphi \rightarrow \varphi$ PL, 1, 2
4. **KDB4** $\vdash \Box\varphi \rightarrow \Box\Box\varphi$ 4
5. **KDB4** $\vdash \Box\varphi \rightarrow \varphi$ PL, 4, 3.

4. **KB4** $\vdash$ 5:

1. **KB4** $\vdash \Diamond\varphi \rightarrow \Box\Diamond\Diamond\varphi$ B with $\Diamond\varphi$ for p
2. **KB4** $\vdash \Diamond\Diamond\varphi \rightarrow \Diamond\varphi$ $4_\Diamond$
3. **KB4** $\vdash \Box\Diamond\Diamond\varphi \rightarrow \Box\Diamond\varphi$ RK, 2
4. **KB4** $\vdash \Diamond\varphi \rightarrow \Box\Diamond\varphi$ PL, 1, 3.

5. **KB5** $\vdash$ 4:

1. **KB5** $\vdash \Box\varphi \rightarrow \Box\Diamond\Box\varphi$ B with $\Box\varphi$ for p
2. **KB5** $\vdash \Diamond\Box\varphi \rightarrow \Box\varphi$ $5_\Diamond$
3. **KB5** $\vdash \Box\Diamond\Box\varphi \rightarrow \Box\Box\varphi$ RK, 2
4. **KB5** $\vdash \Box\varphi \rightarrow \Box\Box\varphi$ PL, 1, 3.

6. **KT** $\vdash$ D:

1. **KT** $\vdash \Box\varphi \rightarrow \varphi$ T
2. **KT** $\vdash \varphi \rightarrow \Diamond\varphi$ $T_\Diamond$
3. **KT** $\vdash \Box\varphi \rightarrow \Diamond\varphi$ PL, 1, 2

□

Definição prf.2. Following tradition, we define **S4** to be the system **KT4**, and **S5** the system **KT5**.

The following proposition shows that the classical system **S5** has several equivalent axiomatizations. This should not surprise, as the various combinations of axioms all characterize equivalence relations (see ??).

Proposição prf.3. **KT5** = **KTB4** = **KDB4** = **KDB5**.

*nml:prf:prs:
prop:S5*

Prova. Exercise.

□

Problema prf.1. Prove **Proposição prf.3**.

Créditos das Imagens

Referências Bibliográficas