

-NoValue-

tab.1 Soundness for Additional Rules

nml:tab:msn: sec: We say a rule is sound for a class of models if, whenever a branch in a **tableau** is satisfiable in a model from that class, the branch resulting from applying the rule is also satisfiable in a model from that class.

nml:tab:msn: prop:soundness-T **Proposição tab.1.** $T\Box$ and $T\Diamond$ are sound for reflexive models.

Prova. 1. The branch is expanded by applying $T\Box$ to $\sigma T\Box\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma T\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma) \Vdash \Box\psi$. Since R is reflexive, we know that $Rf(\sigma)f(\sigma)$. Hence, $\mathfrak{M}, f(\sigma) \Vdash \psi$, O.s., $\mathfrak{M}, f$ satisfies $\sigma T\psi$.

2. The branch is expanded by applying $T\Diamond$ to $\sigma F\Diamond\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma F\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma) \not\Vdash \Diamond\psi$. Since R is reflexive, we know that $Rf(\sigma)f(\sigma)$. Hence, $\mathfrak{M}, f(\sigma) \not\Vdash \psi$, O.s., $\mathfrak{M}, f$ satisfies $\sigma F\psi$. $\square$

nml:tab:msn: prop:soundness-D **Proposição tab.2.** $D\Box$ and $D\Diamond$ are sound for serial models.

Prova. 1. The branch is expanded by applying $D\Box$ to $\sigma T\Box\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma T\Diamond\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma) \Vdash \Box\psi$. Since R is serial, there is a $w \in W$ such that $Rf(\sigma)w$. Then $\mathfrak{M}, w \Vdash \psi$, and hence $\mathfrak{M}, f(\sigma) \Vdash \Diamond\psi$. So, $\mathfrak{M}, f$ satisfies $\sigma T\Diamond\psi$.

2. The branch is expanded by applying $D\Diamond$ to $\sigma F\Diamond\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma F\Box\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma) \not\Vdash \Diamond\psi$. Since R is serial, there is a $w \in W$ such that $Rf(\sigma)w$. Then $\mathfrak{M}, w \not\Vdash \psi$, and hence $\mathfrak{M}, f(\sigma) \not\Vdash \Box\psi$. So, $\mathfrak{M}, f$ satisfies $\sigma F\Box\psi$. $\square$

nml:tab:msn: prop:soundness-B **Proposição tab.3.** $B\Box$ and $B\Diamond$ are sound for symmetric models.

Prova. 1. The branch is expanded by applying $B\Box$ to $\sigma.n T\Box\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma T\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma.n) \Vdash \Box\psi$. Since f is an interpretation of prefixes on the branch into $\mathfrak{M}$, we know that $Rf(\sigma)f(\sigma.n)$. Since R is symmetric, $Rf(\sigma.n)f(\sigma)$. Since $\mathfrak{M}, f(\sigma.n) \Vdash \Box\psi$, $\mathfrak{M}, f(\sigma) \Vdash \psi$. Hence, $\mathfrak{M}, f$ satisfies $\sigma T\psi$.

2. The branch is expanded by applying $B\Diamond$ to $\sigma.n F\Diamond\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma F\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma.n) \not\Vdash \Diamond\psi$. Since f is an interpretation of prefixes on

the branch into $\mathfrak{M}$, we know that $Rf(\sigma)f(\sigma.n)$. Since R is symmetric, $Rf(\sigma.n)f(\sigma)$. Since $\mathfrak{M}, f(\sigma.n) \not\models \Diamond\psi$, $\mathfrak{M}, f(\sigma) \not\models \psi$. Hence, $\mathfrak{M}, f$ satisfies $\sigma \mathbb{F} \psi$. $\square$

Proposição tab.4. $4\Box$ and $4\Diamond$ are sound for transitive models.

*nml:tab:msn:
prop:soundness-4*

- Prova.* 1. The branch is expanded by applying $4\Box$ to $\sigma \mathbb{T} \Box\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma.n \mathbb{T} \Box\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma) \Vdash \Box\psi$. Since f is an interpretation of prefixes on the branch into $\mathfrak{M}$ and $\sigma.n$ must be used, we know that $Rf(\sigma)f(\sigma.n)$. Now let w be any world such that $Rf(\sigma.n)w$. Since R is transitive, $Rf(\sigma)w$. Since $\mathfrak{M}, f(\sigma) \Vdash \Box\psi$, $\mathfrak{M}, w \Vdash \psi$. Hence, $\mathfrak{M}, f(\sigma.n) \Vdash \Box\psi$, and $\mathfrak{M}, f$ satisfies $\sigma.n \mathbb{T} \Box\psi$.
2. The branch is expanded by applying $4\Diamond$ to $\sigma \mathbb{F} \Diamond\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma.n \mathbb{F} \Diamond\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma) \not\models \Diamond\psi$. Since f is an interpretation of prefixes on the branch into $\mathfrak{M}$ and $\sigma.n$ must be used, we know that $Rf(\sigma)f(\sigma.n)$. Now let w be any world such that $Rf(\sigma.n)w$. Since R is transitive, $Rf(\sigma)w$. Since $\mathfrak{M}, f(\sigma) \not\models \Diamond\psi$, $\mathfrak{M}, w \not\models \psi$. Hence, $\mathfrak{M}, f(\sigma.n) \not\models \Diamond\psi$, and $\mathfrak{M}, f$ satisfies $\sigma.n \mathbb{F} \Diamond\psi$. $\square$

Proposição tab.5. $4r\Box$ and $4r\Diamond$ are sound for euclidean models.

*nml:tab:msn:
prop:soundness-4r*

- Prova.* 1. The branch is expanded by applying $4r\Box$ to $\sigma.n \mathbb{T} \Box\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma \mathbb{T} \Box\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma.n) \Vdash \Box\psi$. Since f is an interpretation of prefixes on the branch into $\mathfrak{M}$, we know that $Rf(\sigma)f(\sigma.n)$. Now let w be any world such that $Rf(\sigma)w$. Since R is euclidean, $Rf(\sigma.n)w$. Since $\mathfrak{M}, f(\sigma.n) \Vdash \Box\psi$, $\mathfrak{M}, w \Vdash \psi$. Hence, $\mathfrak{M}, f(\sigma) \Vdash \Box\psi$, and $\mathfrak{M}, f$ satisfies $\sigma \mathbb{T} \Box\psi$.
2. The branch is expanded by applying $4r\Diamond$ to $\sigma.n \mathbb{F} \Diamond\psi \in \Gamma$: This results in a new **fórmula assinada** $\sigma \mathbb{F} \Diamond\psi$ on the branch. Suppose $\mathfrak{M}, f \Vdash \Gamma$, in particular, $\mathfrak{M}, f(\sigma.n) \not\models \Diamond\psi$. Since f is an interpretation of prefixes on the branch into $\mathfrak{M}$, we know that $Rf(\sigma)f(\sigma.n)$. Now let w be any world such that $Rf(\sigma)w$. Since R is euclidean, $Rf(\sigma.n)w$. Since $\mathfrak{M}, f(\sigma.n) \not\models \Diamond\psi$, $\mathfrak{M}, w \not\models \psi$. Hence, $\mathfrak{M}, f(\sigma) \not\models \Diamond\psi$, and $\mathfrak{M}, f$ satisfies $\sigma \mathbb{F} \Diamond\psi$. $\square$

Corolário tab.6. The **tableau** systems given in ?? are sound for the respective classes of models.

*nml:tab:msn:
cor:soundness-logics*

Créditos das Imagens

Referências Bibliográficas