

Capítulo udf

Syntax and Semantics

This is a very quick summary of definitions only. It should be expanded to provide a gentle intro to proofs by induction on formulas, with lots more examples.

-NoValue-

syn.1 Introduction

pl:syn:int:
sec

Propositional logic deals with **fórmulas** that are built from **variáveis proposicionais** using the propositional connectives $\neg$, $\wedge$, $\vee$, $\rightarrow$, and $\leftrightarrow$. Intuitively, a **variável proposicional** p stands for a sentence or proposition that is true or false. Whenever the “truth value” of the **variável proposicional** in a **fórmula** is determined, so is the truth value of any **fórmulas** formed from them using propositional connectives. We say that propositional logic is *truth functional*, because its semantics is given by functions of truth values. In particular, in propositional logic we leave out of consideration any further determination of truth and falsity, p.ex., whether something is necessarily true rather than just contingently true, or whether something is known to be true, or whether something is true now rather than was true or will be true. We only consider two truth values true ($\mathbb{T}$) and false ($\mathbb{F}$), and so exclude from discussion the possibility that a statement may be neither true nor false, or only half true. We also concentrate only on connectives where the truth value of a **fórmula** built from them is completely determined by the truth values of its parts (and not, say, on its meaning). In particular, whether the truth value of conditionals in English is truth functional in this sense is contentious. The material conditional $\rightarrow$ is; other logics deal with conditionals that are not truth functional.

In order to develop the theory and metatheory of truth-functional propositional logic, we must first define the syntax and semantics of its expressions. We will describe one way of constructing **fórmulas** from **variáveis proposicio-**

naís using the connectives. Alternative definitions are possible. Other systems will choose different symbols, will select different sets of connectives as primitive, and will use parentheses differently (or even not at all, as in the case of so-called Polish notation). What all approaches have in common, though, is that the formation rules define the set of *fórmulas* *inductively*. If done properly, every expression can result essentially in only one way according to the formation rules. The inductive definition resulting in expressions that are *uniquely readable* means we can give meanings to these expressions using the same method—inductive definition.

Giving the meaning of expressions is the domain of semantics. The central concept in semantics for propositional logic is that of satisfaction in a *valoração*. A *valoração* $\mathbf{v}$ assigns truth values $\mathbb{T}, \mathbb{F}$ to the *variáveis proposicionais*. Any *valoração* determines a truth value $\bar{\mathbf{v}}(\varphi)$ for any *fórmula* φ . A *fórmula* is satisfied in a *valoração* $\mathbf{v}$ iff $\bar{\mathbf{v}}(\varphi) = \mathbb{T}$ —we write this as $\mathbf{v} \models \varphi$. This relation can also be defined by induction on the structure of φ , using the truth functions for the logical connectives to define, say, satisfaction of $\varphi \wedge \psi$ in terms of satisfaction (or not) of φ and ψ .

On the basis of the satisfaction relation $\mathbf{v} \models \varphi$ for sentences we can then define the basic semantic notions of tautology, entailment, and satisfiability. A *fórmula* is a tautology, $\models \varphi$, if every *valoração* satisfies it, O.s., $\bar{\mathbf{v}}(\varphi) = \mathbb{T}$ for any $\mathbf{v}$. It is entailed by a set of *fórmulas*, $\Gamma \models \varphi$, if every *valoração* that satisfies all the *fórmulas* in Γ also satisfies φ . And a set of *fórmulas* is satisfiable if some *valoração* satisfies all *fórmulas* in it at the same time. Because *fórmulas* are inductively defined, and satisfaction is in turn defined by induction on the structure of *fórmulas*, we can use induction to prove properties of our semantics and to relate the semantic notions defined.

-NoValue-

syn.2 Propositional Fórmulas

Fórmulas of propositional logic are built up from *variáveis proposicionais*, the propositional constant $\perp$ and the propositional constant $\top$ using *logical connectives*. pl:syn:fml:
sec

1. A *infinito enumerável* set At_0 of *variáveis proposicionais* $p_0, p_1, \dots$
2. The propositional constant for *falsidade* $\perp$.
3. The propositional constant for *verdade* $\top$.
4. The logical connectives: $\neg$ (negation), $\wedge$ (conjunction), $\vee$ (disjunction), $\rightarrow$ (*condicional*), $\leftrightarrow$ (*bicondicional*)
5. Punctuation marks: $(,)$, and the comma.

We denote this language of propositional logic by $\mathcal{L}_0$.

You may be familiar with different terminology and symbols than the ones we use above. Logic texts (and teachers) commonly use either $\sim$, $\neg$, and $!$ for “negation”, $\wedge$, $\cdot$, and $\&$ for “conjunction”. Commonly used symbols for the “conditional” or “implication” are $\rightarrow$, $\Rightarrow$, and $\supset$. Symbols for “biconditional,” “bi-implication,” or “(material) equivalence” are $\leftrightarrow$, $\Leftrightarrow$, and $\equiv$. The $\perp$ symbol is variously called “falsity,” “falsum,” “absurdity,” or “bottom.” The $\top$ symbol is variously called “truth,” “verum,” or “top.”

pl:syn:fml:
defn:formulas

Definição syn.1 (Formula). The set $\text{Frm}(\mathcal{L}_0)$ of *fórmulas* of propositional logic is defined inductively as follows:

1. $\perp$ is an atomic *fórmula*.
2. $\top$ is an atomic *fórmula*.
3. Every *variável proposicional* p_i is an atomic *fórmula*.
4. If φ is a *fórmula*, then $\neg\varphi$ is a *fórmula*.
5. If φ and ψ are *fórmulas*, then $(\varphi \wedge \psi)$ is a *fórmula*.
6. If φ and ψ are *fórmulas*, then $(\varphi \vee \psi)$ is a *fórmula*.
7. If φ and ψ are *fórmulas*, then $(\varphi \rightarrow \psi)$ is a *fórmula*.
8. If φ and ψ are *fórmulas*, then $(\varphi \leftrightarrow \psi)$ is a *fórmula*.
9. Nothing else is a *fórmula*.

The definition of *fórmulas* is an *inductive definition*. Essentially, we construct the set of *fórmulas* in infinitely many stages. In the initial stage, we pronounce all atomic formulas to be formulas; this corresponds to the first few cases of the definition, O.s., the cases for $\top$, $\perp$, p_i . “Atomic *fórmula*” thus means any *fórmula* of this form.

The other cases of the definition give rules for constructing new *fórmulas* out of *fórmulas* already constructed. At the second stage, we can use them to construct *fórmulas* out of atomic *fórmulas*. At the third stage, we construct new formulas from the atomic formulas and those obtained in the second stage, and so on. A *fórmula* is anything that is eventually constructed at such a stage, and nothing else.

When writing a formula $(\psi * \chi)$ constructed from ψ , χ using a two-place connective $*$, we will often leave out the outermost pair of parentheses and write simply $\psi * \chi$.

Definição syn.2 (Syntactic identity). The symbol $\equiv$ expresses syntactic identity between strings of symbols, O.s., $\varphi \equiv \psi$ iff φ and ψ are strings of symbols of the same length and which contain the same symbol in each place.

The $\equiv$ symbol may be flanked by strings obtained by concatenation, p.ex., $\varphi \equiv (\psi \vee \chi)$ means: the string of symbols φ is the same string as the one obtained by concatenating an opening parenthesis, the string ψ , the $\vee$ symbol, the string χ , and a closing parenthesis, in this order. If this is the case, then we know that the first symbol of φ is an opening parenthesis, φ contains ψ as a substring (starting at the second symbol), that substring is followed by $\vee$, etc.

-NoValue-

syn.3 Preliminaries

Teorema syn.3 (Principle of induction on fórmulas). *If some property P holds for all the atomic fórmulas and is such that*

pl:syn:pre:
P^{sc}
pl:syn:pre:
thm:induction

1. *it holds for $\neg\varphi$ whenever it holds for φ ;*
2. *it holds for $(\varphi \wedge \psi)$ whenever it holds for φ and ψ ;*
3. *it holds for $(\varphi \vee \psi)$ whenever it holds for φ and ψ ;*
4. *it holds for $(\varphi \rightarrow \psi)$ whenever it holds for φ and ψ ;*
5. *it holds for $(\varphi \leftrightarrow \psi)$ whenever it holds for φ and ψ ;*

then P holds for all fórmulas.

Prova. Let S be the collection of all fórmulas with property P . Clearly $S \subseteq \text{Frm}(\mathcal{L}_0)$. S satisfies all the conditions of Definição syn.1: it contains all atomic fórmulas and is closed under the operadores lógicos. $\text{Frm}(\mathcal{L}_0)$ is the smallest such class, so $\text{Frm}(\mathcal{L}_0) \subseteq S$. So $\text{Frm}(\mathcal{L}_0) = S$, and every formula has property P . $\square$

Proposição syn.4. *Any fórmula in $\text{Frm}(\mathcal{L}_0)$ is balanced, in that it has as many left parentheses as right ones.*

pl:syn:pre:
prop:balanced

Problema syn.1. Prove Proposição syn.4

Proposição syn.5. *No proper initial segment of a fórmula is a fórmula.*

pl:syn:pre:
prop:noinit

Problema syn.2. Prove Proposição syn.5

Proposição syn.6 (Unique Readability). *Any fórmula φ in $\text{Frm}(\mathcal{L}_0)$ has exactly one parsing as one of the following*

1. $\perp$.
2. $\top$.
3. p_n for some $p_n \in \text{At}_0$.

4. $\neg\psi$ for some *fórmula* ψ .
5. $(\psi \wedge \chi)$ for some *fórmulas* ψ and χ .
6. $(\psi \vee \chi)$ for some *fórmulas* ψ and χ .
7. $(\psi \rightarrow \chi)$ for some *fórmulas* ψ and χ .
8. $(\psi \leftrightarrow \chi)$ for some *fórmulas* ψ and χ .

Moreover, this parsing is unique.

Prova. By induction on φ . For instance, suppose that φ has two distinct readings as $(\psi \rightarrow \chi)$ and $(\psi' \rightarrow \chi')$. Then ψ and ψ' must be the same (or else one would be a proper initial segment of the other); so if the two readings of φ are distinct it must be because χ and χ' are distinct readings of the same sequence of symbols, which is impossible by the inductive hypothesis. $\square$

Definição syn.7 (Uniform Substitution). If φ and ψ are *fórmulas*, and p_i is a *variável proposicional*, then $\varphi[\psi/p_i]$ denotes the result of replacing each occurrence of p_i by an occurrence of ψ in φ ; similarly, the simultaneous substitution of $p_1, \dots, p_n$ by *fórmulas* $\psi_1, \dots, \psi_n$ is denoted by $\varphi[\psi_1/p_1, \dots, \psi_n/p_n]$.

Problema syn.3. For each of the five *fórmulas* below determine whether the *fórmula* can be expressed as a substitution $\varphi[\psi/p_i]$ where φ is (i) p_0 ; (ii) $(\neg p_0 \wedge p_1)$; and (iii) $((\neg p_0 \rightarrow p_1) \wedge p_2)$. In each case specify the relevant substitution.

1. p_1
2. $(\neg p_0 \wedge p_0)$
3. $((p_0 \vee p_1) \wedge p_2)$
4. $\neg((p_0 \rightarrow p_1) \wedge p_2)$
5. $((\neg(p_0 \rightarrow p_1) \rightarrow (p_0 \vee p_1)) \wedge \neg(p_0 \wedge p_1))$

Problema syn.4. Give a mathematically rigorous definition of $\varphi[\psi/p]$ by induction.

-NoValue-

syn.4 Formation Sequences

Defining *fórmulas* via an inductive definition, and the complementary technique of proving properties of *fórmulas* via induction, is an elegant and efficient approach. However, it can also be useful to consider a more bottom-up, step-by-step approach to the construction of *fórmulas*, which we do here using the notion of a *formation sequence*.

Definição syn.8 (Formation sequences for formulas). A finite sequence $\langle \varphi_0, \dots, \varphi_n \rangle$ of strings of symbols from the language $\mathcal{L}_0$ is a *formation sequence* for φ if $\varphi \equiv \varphi_n$ and for all $i \leq n$, either φ_i is an atomic formula or there exist $j, k < i$ such that one of the following holds:

pl:syn:fseq:
defn:fseq-frn

1. $\varphi_i \equiv \neg \varphi_j$.
2. $\varphi_i \equiv (\varphi_j \wedge \varphi_k)$.
3. $\varphi_i \equiv (\varphi_j \vee \varphi_k)$.
4. $\varphi_i \equiv (\varphi_j \rightarrow \varphi_k)$.
5. $\varphi_i \equiv (\varphi_j \leftrightarrow \varphi_k)$.

Exemplo syn.9.

$$\langle p_0, p_1, (p_1 \wedge p_0), \neg(p_1 \wedge p_0) \rangle$$

is a formation sequence of $\neg(p_1 \wedge p_0)$, as is

$$\langle p_0, p_1, p_0, (p_1 \wedge p_0), (p_0 \rightarrow p_1), \neg(p_1 \wedge p_0) \rangle.$$

As can be seen from the second example, formation sequences may contain ‘junk’: formulas which are redundant or do not contribute to the construction.

Proposição syn.10. Every *fórmula* φ in $\text{Frm}(\mathcal{L}_0)$ has a formation sequence.

pl:syn:fseq:
prop:formed

Prova. Suppose φ is atomic. Then the sequence $\langle \varphi \rangle$ is a formation sequence for φ . Now suppose that ψ and χ have formation sequences $\langle \psi_0, \dots, \psi_n \rangle$ and $\langle \chi_0, \dots, \chi_m \rangle$ respectively.

1. If $\varphi \equiv \neg \psi$, then $\langle \psi_0, \dots, \psi_n, \neg \psi_n \rangle$ is a formation sequence for φ .
2. If $\varphi \equiv (\psi \wedge \chi)$, then $\langle \psi_0, \dots, \psi_n, \chi_0, \dots, \chi_m, (\psi_n \wedge \chi_m) \rangle$ is a formation sequence for φ .
3. If $\varphi \equiv (\psi \vee \chi)$, then $\langle \psi_0, \dots, \psi_n, \chi_0, \dots, \chi_m, (\psi_n \vee \chi_m) \rangle$ is a formation sequence for φ .
4. If $\varphi \equiv (\psi \rightarrow \chi)$, then $\langle \psi_0, \dots, \psi_n, \chi_0, \dots, \chi_m, (\psi_n \rightarrow \chi_m) \rangle$ is a formation sequence for φ .
5. If $\varphi \equiv (\psi \leftrightarrow \chi)$, then $\langle \psi_0, \dots, \psi_n, \chi_0, \dots, \chi_m, (\psi_n \leftrightarrow \chi_m) \rangle$ is a formation sequence for φ .

By the principle of induction on *fórmulas*, every *fórmula* has a formation sequence. $\square$

We can also prove the converse. This is important because it shows that our two ways of defining formulas are equivalent: they give the same results. It also means that we can prove theorems about formulas by using ordinary induction on the length of formation sequences.

*pl:syn:fseq:
lem:fseq-init* **Lema syn.11.** Suppose that $\langle \varphi_0, \dots, \varphi_n \rangle$ is a formation sequence for φ_n , and that $k \leq n$. Then $\langle \varphi_0, \dots, \varphi_k \rangle$ is a formation sequence for φ_k .

Prova. Exercise. □

*pl:syn:fseq:
thm:fseq-frm-equiv* **Teorema syn.12.** $\text{Frm}(\mathcal{L}_0)$ is the set of all strings of symbols in the language $\mathcal{L}_0$ with a formation sequence.

Prova. Let F be the set of all strings of symbols in the language $\mathcal{L}_0$ that have a formation sequence. We have seen in **Proposição syn.10** that $\text{Frm}(\mathcal{L}_0) \subseteq F$, so now we prove the converse.

Suppose φ has a formation sequence $\langle \varphi_0, \dots, \varphi_n \rangle$. We prove that $\varphi \in \text{Frm}(\mathcal{L}_0)$ by strong induction on n . Our induction hypothesis is that every string of symbols with a formation sequence of length $m < n$ is in $\text{Frm}(\mathcal{L}_0)$. By the definition of a formation sequence, either φ_n is atomic or there must exist $j, k < n$ such that one of the following is the case:

1. $\varphi_n \equiv \neg \varphi_j$.
2. $\varphi_n \equiv (\varphi_j \wedge \varphi_k)$.
3. $\varphi_n \equiv (\varphi_j \vee \varphi_k)$.
4. $\varphi_n \equiv (\varphi_j \rightarrow \varphi_k)$.
5. $\varphi_n \equiv (\varphi_j \leftrightarrow \varphi_k)$.

Now we reason by cases. If φ_n is atomic then $\varphi_n \in \text{Frm}(\mathcal{L}_0)$. Suppose instead that $\varphi \equiv (\varphi_j \wedge \varphi_k)$. By **Lema syn.11**, $\langle \varphi_0, \dots, \varphi_j \rangle$ and $\langle \varphi_0, \dots, \varphi_k \rangle$ are formation sequences for φ_j and φ_k respectively. Since these are proper initial subsequences of the formation sequence for φ , they both have length less than n . Therefore by the induction hypothesis, φ_j and φ_k are in $\text{Frm}(\mathcal{L}_0)$, and so by the definition of **a fórmula**, so is $(\varphi_j \wedge \varphi_k)$. The other cases follow by parallel reasoning. □

-NoValue-

syn.5 Valorações and Satisfaction

*pl:syn:val:
sec*

Definição syn.13 (Valorações). Let $\{\mathbb{T}, \mathbb{F}\}$ be the set of the two truth values, “true” and “false.” A *valoração* for $\mathcal{L}_0$ is a function $\mathbf{v}$ assigning either $\mathbb{T}$ or $\mathbb{F}$ to the **variáveis proposicionais** of the language, O.s., $\mathbf{v}: \text{At}_0 \rightarrow \{\mathbb{T}, \mathbb{F}\}$.

Definição syn.14. Given a **avalição** $\mathbf{v}$, define the evaluation function $\bar{\mathbf{v}}: \text{Frm}(\mathcal{L}_0) \rightarrow \{\mathbb{T}, \mathbb{F}\}$ inductively by:

$$\begin{aligned}\bar{\mathbf{v}}(\perp) &= \mathbb{F}; \\ \bar{\mathbf{v}}(\top) &= \mathbb{T}; \\ \bar{\mathbf{v}}(p_n) &= \mathbf{v}(p_n); \\ \bar{\mathbf{v}}(\neg\varphi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F}; \\ \mathbb{F} & \text{otherwise.} \end{cases} \\ \bar{\mathbf{v}}(\varphi \wedge \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T} \text{ and } \bar{\mathbf{v}}(\psi) = \mathbb{T}; \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F} \text{ or } \bar{\mathbf{v}}(\psi) = \mathbb{F}. \end{cases} \\ \bar{\mathbf{v}}(\varphi \vee \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T} \text{ or } \bar{\mathbf{v}}(\psi) = \mathbb{T}; \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F} \text{ and } \bar{\mathbf{v}}(\psi) = \mathbb{F}. \end{cases} \\ \bar{\mathbf{v}}(\varphi \rightarrow \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F} \text{ or } \bar{\mathbf{v}}(\psi) = \mathbb{T}; \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T} \text{ and } \bar{\mathbf{v}}(\psi) = \mathbb{F}. \end{cases} \\ \bar{\mathbf{v}}(\varphi \leftrightarrow \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \bar{\mathbf{v}}(\psi); \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) \neq \bar{\mathbf{v}}(\psi). \end{cases}\end{aligned}$$

explanation

The clauses correspond to the following truth tables:

φ	$\neg\varphi$	φ	ψ	$\varphi \wedge \psi$	φ	ψ	$\varphi \vee \psi$
$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$
$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$
$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{T}$
$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$

φ	ψ	$\varphi \rightarrow \psi$	φ	ψ	$\varphi \leftrightarrow \psi$
$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$
$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$
$\mathbb{F}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$
$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$

Problema syn.5. Consider adding to $\mathcal{L}_0$ a ternary connective $\diamond$ with evaluation given by

$$\bar{\mathbf{v}}(\diamond(\varphi, \psi, \chi)) = \begin{cases} \bar{\mathbf{v}}(\psi) & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T}; \\ \bar{\mathbf{v}}(\chi) & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F}. \end{cases}$$

Write down the truth table for this connective.

Teorema syn.15 (Local Determination). Suppose that $\mathbf{v}_1$ and $\mathbf{v}_2$ are **avalições** that agree on the **variáveis proposicionais** occurring in φ , *O.s.*, $\mathbf{v}_1(p_n) = \mathbf{v}_2(p_n)$ whenever p_n occurs in some **fórmula** φ . Then $\bar{\mathbf{v}}_1$ and $\bar{\mathbf{v}}_2$ also agree on φ , *O.s.*, $\bar{\mathbf{v}}_1(\varphi) = \bar{\mathbf{v}}_2(\varphi)$.

Prova. By induction on φ . □

pl:syn:val:
defn:satisfaction

Definição syn.16 (Satisfaction). We can inductively define the notion of *satisfaction of a fórmula* φ by a *valoração* $\mathbf{v}$, $\mathbf{v} \models \varphi$, as follows. (We write $\mathbf{v} \not\models \varphi$ to mean “not $\mathbf{v} \models \varphi$.”)

1. $\varphi \equiv \perp$: $\mathbf{v} \not\models \varphi$.
2. $\varphi \equiv \top$: $\mathbf{v} \models \varphi$.
3. $\varphi \equiv p_i$: $\mathbf{v} \models \varphi$ iff $\mathbf{v}(p_i) = \mathbb{T}$.
4. $\varphi \equiv \neg\psi$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \not\models \psi$.
5. $\varphi \equiv (\psi \wedge \chi)$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \models \psi$ and $\mathbf{v} \models \chi$.
6. $\varphi \equiv (\psi \vee \chi)$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \models \psi$ or $\mathbf{v} \models \chi$ (or both).
7. $\varphi \equiv (\psi \rightarrow \chi)$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \not\models \psi$ or $\mathbf{v} \models \chi$ (or both).
8. $\varphi \equiv (\psi \leftrightarrow \chi)$: $\mathbf{v} \models \varphi$ iff either both $\mathbf{v} \models \psi$ and $\mathbf{v} \models \chi$, or neither $\mathbf{v} \models \psi$ nor $\mathbf{v} \models \chi$.

If Γ is a set of *fórmulas*, $\mathbf{v} \models \Gamma$ iff $\mathbf{v} \models \varphi$ for every $\varphi \in \Gamma$.

pl:syn:val:
prop:sat-value

Proposição syn.17. $\mathbf{v} \models \varphi$ iff $\overline{\mathbf{v}}(\varphi) = \mathbb{T}$.

Prova. By induction on φ . □

Problema syn.6. Prove **Proposição syn.17**

-NoValue-

syn.6 Semantic Notions

pl:syn:sem:
sec

We define the following semantic notions:

- Definição syn.18.**
1. A *fórmula* φ is *satisfiable* if for some $\mathbf{v}$, $\mathbf{v} \models \varphi$; it is *unsatisfiable* if for no $\mathbf{v}$, $\mathbf{v} \models \varphi$;
 2. A *fórmula* φ is a *tautology* if $\mathbf{v} \models \varphi$ for all *valorações* $\mathbf{v}$;
 3. A *fórmula* φ is *contingent* if it is satisfiable but not a tautology;
 4. If Γ is a set of *fórmulas*, $\Gamma \models \varphi$ (“ Γ entails φ ”) if and only if $\mathbf{v} \models \varphi$ for every *valoração* $\mathbf{v}$ for which $\mathbf{v} \models \Gamma$.
 5. If Γ is a set of *fórmulas*, Γ is *satisfiable* if there is a *valoração* $\mathbf{v}$ for which $\mathbf{v} \models \Gamma$, and Γ is *unsatisfiable* otherwise.

Problema syn.7. For each of the following four *fórmulas* determine whether it is (a) satisfiable, (b) tautology, and (c) contingent.

1. $(p_0 \rightarrow (\neg p_1 \rightarrow \neg p_0))$.
2. $((p_0 \wedge \neg p_1) \rightarrow (\neg p_0 \wedge p_2)) \leftrightarrow ((p_2 \rightarrow p_0) \rightarrow (p_0 \rightarrow p_1))$.
3. $(p_0 \leftrightarrow p_1) \rightarrow (p_2 \leftrightarrow \neg p_1)$.
4. $((p_0 \leftrightarrow (\neg p_1 \wedge p_2)) \vee (p_2 \rightarrow (p_0 \leftrightarrow p_1)))$.

Proposição syn.19.

*pl:syn:sem:
prop:semanticalfacts*

1. φ is a tautology if and only if $\emptyset \models \varphi$;
2. If $\Gamma \models \varphi$ and $\Gamma \models \varphi \rightarrow \psi$ then $\Gamma \models \psi$;
3. If Γ is satisfiable then every finite subset of Γ is also satisfiable;
4. Monotonicity: if $\Gamma \subseteq \Delta$ and $\Gamma \models \varphi$ then also $\Delta \models \varphi$;
5. Transitivity: if $\Gamma \models \varphi$ and $\Delta \cup \{\varphi\} \models \psi$ then $\Gamma \cup \Delta \models \psi$.

*pl:syn:sem:
def:monotonicity
pl:syn:sem:
def:Cut*

Prova. Exercise. □

Problema syn.8. Prove **Proposição syn.19**

Proposição syn.20. $\Gamma \models \varphi$ if and only if $\Gamma \cup \{\neg\varphi\}$ is unsatisfiable.

*pl:syn:sem:
prop:entails-unsat*

Prova. Exercise. □

Problema syn.9. Prove **Proposição syn.20**

Teorema syn.21 (Semantic Deduction Theorem). $\Gamma \models \varphi \rightarrow \psi$ if and only if $\Gamma \cup \{\varphi\} \models \psi$.

*pl:syn:sem:
thm:sem-deduction*

Prova. Exercise. □

Problema syn.10. Prove **Teorema syn.21**

Créditos das Imagens

Referências Bibliográficas