

-NoValue-

syn.1 Valorações and Satisfaction

pl:syn:val:
sec

Definição syn.1 (Valorações). Let $\{\mathbb{T}, \mathbb{F}\}$ be the set of the two truth values, “true” and “false.” A *valoração* for $\mathcal{L}_0$ is a function $\mathbf{v}$ assigning either $\mathbb{T}$ or $\mathbb{F}$ to the *variáveis proposicionais* of the language, O.s., $\mathbf{v}: \text{At}_0 \rightarrow \{\mathbb{T}, \mathbb{F}\}$.

Definição syn.2. Given a *valoração* $\mathbf{v}$, define the evaluation function $\bar{\mathbf{v}}: \text{Frm}(\mathcal{L}_0) \rightarrow \{\mathbb{T}, \mathbb{F}\}$ inductively by:

$$\begin{aligned} \bar{\mathbf{v}}(\perp) &= \mathbb{F}; \\ \bar{\mathbf{v}}(\top) &= \mathbb{T}; \\ \bar{\mathbf{v}}(\rho_n) &= \mathbf{v}(\rho_n); \\ \bar{\mathbf{v}}(\neg\varphi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F}; \\ \mathbb{F} & \text{otherwise.} \end{cases} \\ \bar{\mathbf{v}}(\varphi \wedge \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T} \text{ and } \bar{\mathbf{v}}(\psi) = \mathbb{T}; \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F} \text{ or } \bar{\mathbf{v}}(\psi) = \mathbb{F}. \end{cases} \\ \bar{\mathbf{v}}(\varphi \vee \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T} \text{ or } \bar{\mathbf{v}}(\psi) = \mathbb{T}; \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F} \text{ and } \bar{\mathbf{v}}(\psi) = \mathbb{F}. \end{cases} \\ \bar{\mathbf{v}}(\varphi \rightarrow \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F} \text{ or } \bar{\mathbf{v}}(\psi) = \mathbb{T}; \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T} \text{ and } \bar{\mathbf{v}}(\psi) = \mathbb{F}. \end{cases} \\ \bar{\mathbf{v}}(\varphi \leftrightarrow \psi) &= \begin{cases} \mathbb{T} & \text{if } \bar{\mathbf{v}}(\varphi) = \bar{\mathbf{v}}(\psi); \\ \mathbb{F} & \text{if } \bar{\mathbf{v}}(\varphi) \neq \bar{\mathbf{v}}(\psi). \end{cases} \end{aligned}$$

The clauses correspond to the following truth tables:

explanation

φ	$\neg\varphi$	φ	ψ	$\varphi \wedge \psi$	φ	ψ	$\varphi \vee \psi$
$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$
$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$
$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{T}$
$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{F}$

φ	ψ	$\varphi \rightarrow \psi$	φ	ψ	$\varphi \leftrightarrow \psi$
$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{T}$
$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$
$\mathbb{F}$	$\mathbb{T}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$
$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$	$\mathbb{F}$	$\mathbb{F}$	$\mathbb{T}$

Problema syn.1. Consider adding to $\mathcal{L}_0$ a ternary connective $\Diamond$ with evaluation given by

$$\bar{\mathbf{v}}(\Diamond(\varphi, \psi, \chi)) = \begin{cases} \bar{\mathbf{v}}(\psi) & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{T}; \\ \bar{\mathbf{v}}(\chi) & \text{if } \bar{\mathbf{v}}(\varphi) = \mathbb{F}. \end{cases}$$

Write down the truth table for this connective.

Teorema syn.3 (Local Determination). Suppose that $\mathbf{v}_1$ and $\mathbf{v}_2$ are *valorações* that agree on the *variáveis proposicionais* occurring in φ , O.s., $\mathbf{v}_1(p_n) = \mathbf{v}_2(p_n)$ whenever p_n occurs in some *fórmula* φ . Then $\bar{\mathbf{v}}_1$ and $\bar{\mathbf{v}}_2$ also agree on φ , O.s., $\bar{\mathbf{v}}_1(\varphi) = \bar{\mathbf{v}}_2(\varphi)$. pl:syn:val:
thm:LocalDetermination

Prova. By induction on φ . □

Definição syn.4 (Satisfaction). We can inductively define the notion of *satisfaction of a fórmula φ by a valoração $\mathbf{v}$* , $\mathbf{v} \models \varphi$, as follows. (We write $\mathbf{v} \not\models \varphi$ to mean “not $\mathbf{v} \models \varphi$.”) pl:syn:val:
defn:satisfaction

1. $\varphi \equiv \perp$: $\mathbf{v} \not\models \varphi$.
2. $\varphi \equiv \top$: $\mathbf{v} \models \varphi$.
3. $\varphi \equiv p_i$: $\mathbf{v} \models \varphi$ iff $\mathbf{v}(p_i) = \mathbb{T}$.
4. $\varphi \equiv \neg\psi$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \not\models \psi$.
5. $\varphi \equiv (\psi \wedge \chi)$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \models \psi$ and $\mathbf{v} \models \chi$.
6. $\varphi \equiv (\psi \vee \chi)$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \models \psi$ or $\mathbf{v} \models \chi$ (or both).
7. $\varphi \equiv (\psi \rightarrow \chi)$: $\mathbf{v} \models \varphi$ iff $\mathbf{v} \not\models \psi$ or $\mathbf{v} \models \chi$ (or both).
8. $\varphi \equiv (\psi \leftrightarrow \chi)$: $\mathbf{v} \models \varphi$ iff either both $\mathbf{v} \models \psi$ and $\mathbf{v} \models \chi$, or neither $\mathbf{v} \models \psi$ nor $\mathbf{v} \models \chi$.

If Γ is a set of *fórmulas*, $\mathbf{v} \models \Gamma$ iff $\mathbf{v} \models \varphi$ for every $\varphi \in \Gamma$.

Proposição syn.5. $\mathbf{v} \models \varphi$ iff $\bar{\mathbf{v}}(\varphi) = \mathbb{T}$. pl:syn:val:
prop:sat-value

Prova. By induction on φ . □

Problema syn.2. Prove **Proposição syn.5**

Créditos das Imagens

Referências Bibliográficas