

-NoValue-

met.1 Second-order Arithmetic

sol:met:spa:
sec Recall that the theory **PA** of Peano arithmetic includes the eight axioms of **Q**,

$$\begin{aligned} & \forall x \, x' \neq 0 \\ & \forall x \, \forall y \, (x' = y' \rightarrow x = y) \\ & \forall x \, (x = 0 \vee \exists y \, x = y') \\ & \forall x \, (x + 0) = x \\ & \forall x \, \forall y \, (x + y') = (x + y)' \\ & \forall x \, (x \times 0) = 0 \\ & \forall x \, \forall y \, (x \times y') = ((x \times y) + x) \\ & \forall x \, \forall y \, (x < y \leftrightarrow \exists z \, (z' + x) = y) \end{aligned}$$

plus all sentences of the form

$$(\varphi(0) \wedge \forall x \, (\varphi(x) \rightarrow \varphi(x'))) \rightarrow \forall x \, \varphi(x).$$

The latter is a “schema,” O.s., a pattern that generates infinitely many **sentenças** of the language of arithmetic, one for each **fórmula** $\varphi(x)$. We call this schema the (first-order) *axiom schema of induction*. In *second-order* Peano arithmetic **PA²**, induction can be stated as a single sentence. **PA²** consists of the first eight axioms above plus the (second-order) *induction axiom*:

$$\forall X \, ((X(0) \wedge \forall x \, (X(x) \rightarrow X(x'))) \rightarrow \forall x \, X(x)).$$

It says that if a subset X of the **domínio** contains $0^{\mathfrak{M}}$ and with any $x \in |\mathfrak{M}|$ also contains x' (O.s., it is “closed under successor”) it contains everything in the **domínio** (O.s., $X = |\mathfrak{M}|$).

The induction axiom guarantees that any **estrutura** satisfying it contains only those **membros** of $|\mathfrak{M}|$ the axioms require to be there, O.s., the values of $\bar{n}$ for $n \in \mathbb{N}$. A model of **PA²** contains no non-standard numbers.

sol:met:spa:
thm:sol-pa-standard **Teorema met.1.** *If $\mathfrak{M} \models \mathbf{PA}^2$ then $|\mathfrak{M}| = \{\text{Val}^{\mathfrak{M}}(\bar{n}) : n \in \mathbb{N}\}$.*

Prova. Let $N = \{\text{Val}^{\mathfrak{M}}(\bar{n}) : n \in \mathbb{N}\}$, and suppose $\mathfrak{M} \models \mathbf{PA}^2$. Of course, for any $n \in \mathbb{N}$, $\text{Val}^{\mathfrak{M}}(\bar{n}) \in |\mathfrak{M}|$, so $N \subseteq |\mathfrak{M}|$.

Now for inclusion in the other direction. Consider a variable assignment s with $s(X) = N$. By assumption,

$$\begin{aligned} \mathfrak{M} & \models \forall X \, ((X(0) \wedge \forall x \, (X(x) \rightarrow X(x'))) \rightarrow \forall x \, X(x)), \text{ thus} \\ \mathfrak{M}, s & \models (X(0) \wedge \forall x \, (X(x) \rightarrow X(x'))) \rightarrow \forall x \, X(x). \end{aligned}$$

Consider the antecedent of this conditional. $\text{Val}^{\mathfrak{M}}(0) \in N$, and so $\mathfrak{M}, s \models X(0)$. The second conjunct, $\forall x \, (X(x) \rightarrow X(x'))$ is also satisfied. For suppose

$x \in N$. By definition of N , $x = \text{Val}^{\mathfrak{M}}(\bar{n})$ for some n . That gives $\iota^{\mathfrak{M}}(x) = \text{Val}^{\mathfrak{M}}(\bar{n} + \bar{1}) \in N$. So, $\iota^{\mathfrak{M}}(x) \in N$.

We have that $\mathfrak{M}, s \models X(0) \wedge \forall x (X(x) \rightarrow X(x'))$. Consequently, $\mathfrak{M}, s \models \forall x X(x)$. But that means that for every $x \in |\mathfrak{M}|$ we have $x \in s(X) = N$. So, $|\mathfrak{M}| \subseteq N$. $\square$

Corolário met.2. *Any two models of $\mathbf{PA}^2$ are isomorphic.*

*sol:met:spa:
cor:sol-pa-categorical*

Prova. By **Teorema met.1**, the domain of any model of $\mathbf{PA}^2$ is exhausted by $\text{Val}^{\mathfrak{M}}(\bar{n})$. Any such model is also a model of $\mathbf{Q}$. By ??, any such model is standard, O.s., isomorphic to $\mathfrak{N}$. $\square$

Above we defined $\mathbf{PA}^2$ as the theory that contains the first eight arithmetical axioms plus the second-order induction axiom. In fact, thanks to the expressive power of second-order logic, only the *first two* of the arithmetical axioms plus induction are needed for second-order Peano arithmetic.

Proposição met.3. *Let $\mathbf{PA}^{2\ddagger}$ be the second-order theory containing the first two arithmetical axioms (the successor axioms) and the second-order induction axiom. Then $\leq$, $+$, and $\times$ are definable in $\mathbf{PA}^{2\ddagger}$.*

*sol:met:spa:
prop:sol-pa-definable*

Prova. To show that $\leq$ is definable, we have to find a formula $\varphi_{\leq}(x, y)$ such that $\mathfrak{N} \models \varphi_{\leq}(\bar{n}, \bar{m})$ iff $n \leq m$. Consider the formula

$$\psi(x, Y) \equiv Y(x) \wedge \forall y (Y(y) \rightarrow Y(y'))$$

Clearly, $\psi(\bar{n}, Y)$ is satisfied by a set $Y \subseteq \mathbb{N}$ iff $\{m : n \leq m\} \subseteq Y$, so we can take $\varphi_{\leq}(x, y) \equiv \forall Y (\psi(x, Y) \rightarrow Y(y))$.

To see that addition is definable observe that $k + l = m$ iff there is a function u such that $u(0) = k$, $u(n') = u(n)'$ for all n , and $m = u(l)$. We can use this equivalence to define addition in $\mathbf{PA}^{2\ddagger}$ by the following formula:

$$\varphi_+(x, y, z) \equiv \exists u (u(0) = x \wedge \forall w u(x') = u(x)' \wedge u(y) = z)$$

It should be clear that $\mathfrak{N} \models \varphi_+(\bar{k}, \bar{l}, \bar{m})$ iff $k + l = m$. $\square$

Problema met.1. Complete the proof of **Proposição met.3**.

Créditos das Imagens

Referências Bibliográficas