

-NoValue-

syn.1 Describing Infinite and Enumerável Domínios

sol:syn:siz:
sec

A set M is (Dedekind) infinite iff there is uma **injetiva** function $f: M \rightarrow M$ which is not **sobrejetiva**, O.s., with $\text{Im}(f) \neq M$. In first-order logic, we can consider a one-place **símbolo de função** f and say that the function $f^{\mathfrak{M}}$ assigned to it in a **estrutura** $\mathfrak{M}$ is **injetiva** and $\text{Im}(f) \neq |\mathfrak{M}|$:

$$\forall x \forall y (f(x) = f(y) \rightarrow x = y) \wedge \exists y \forall x y \neq f(x).$$

If $\mathfrak{M}$ satisfies this **sentença**, $f^{\mathfrak{M}}: |\mathfrak{M}| \rightarrow |\mathfrak{M}|$ is **injetiva**, and so $|\mathfrak{M}|$ must be infinite. If $|\mathfrak{M}|$ is infinite, and hence such a function exists, we can let $f^{\mathfrak{M}}$ be that function and $\mathfrak{M}$ will satisfy the **sentença**. However, this requires that our language contains the non-logical symbol f which we use for this purpose. In second-order logic, we can simply say that such a function *exists*. This no-longer requires f , and we obtain the **sentença** in pure second-order logic

$$\text{Inf} \equiv \exists u (\forall x \forall y (u(x) = u(y) \rightarrow x = y) \wedge \exists y \forall x y \neq u(x)).$$

$\mathfrak{M} \models \text{Inf}$ iff $|\mathfrak{M}|$ is infinite. We can then define $\text{Fin} \equiv \neg \text{Inf}$; $\mathfrak{M} \models \text{Fin}$ iff $|\mathfrak{M}|$ is finite. No single **sentença** of pure first-order logic can express that the **domínio** is infinite although an infinite set of them can. There is no set of **sentenças** of pure first-order logic that is satisfied in a **estrutura** iff its domain is finite.

Proposição syn.1. $\mathfrak{M} \models \text{Inf}$ iff $|\mathfrak{M}|$ is infinite.

Prova. $\mathfrak{M} \models \text{Inf}$ iff $\mathfrak{M}, s \models \forall x \forall y (u(x) = u(y) \rightarrow x = y) \wedge \exists y \forall x y \neq u(x)$ for some s . If it does, $s(u)$ is uma **injetiva** function, and some $y \in |\mathfrak{M}|$ is not in the range of $s(u)$. Conversely, if there is uma **injetiva** $f: |\mathfrak{M}| \rightarrow |\mathfrak{M}|$ with $\text{Im}(f) \neq |\mathfrak{M}|$, then $s(u) = f$ is such a variable assignment. $\square$

A set M is **enumerável** if there is an enumeration

$$m_0, m_1, m_2, \dots$$

of its **membros** (without repetitions but possibly finite). Such an enumeration exists iff there is um **membro** $z \in M$ and a function $f: M \rightarrow M$ such that $z, f(z), f(f(z)), \dots$, are all the **membros** of M . For if the enumeration exists, $z = m_0$ and $f(m_k) = m_{k+1}$ (or $f(m_k) = m_k$ if m_k is the last **membro** of the enumeration) are the requisite **membro** and function. On the other hand, if such a z and f exist, then $z, f(z), f(f(z)), \dots$, is an enumeration of M , and M is **enumerável**. We can express the existence of z and f in second-order logic to produce a **sentença** true in a **estrutura** iff the **estrutura** is **enumerável**:

$$\text{Count} \equiv \exists z \exists u \forall X ((X(z) \wedge \forall x (X(x) \rightarrow X(u(x)))) \rightarrow \forall x X(x))$$

Proposição syn.2. $\mathfrak{M} \models \text{Count}$ iff $|\mathfrak{M}|$ is **enumerável**.

Prova. Suppose $|\mathfrak{M}|$ is **enumerável**, and let $m_0, m_1, \dots$, be an enumeration. By removing repetitions we can guarantee that no m_k appears twice. Define $f(m_k) = m_{k+1}$ and let $s(z) = m_0$ and $s(u) = f$. We show that

$$\mathfrak{M}, s \models \forall X ((X(z) \wedge \forall x (X(x) \rightarrow X(u(x)))) \rightarrow \forall x X(x))$$

Suppose $M \subseteq |\mathfrak{M}|$ is arbitrary. Suppose further that $\mathfrak{M}, s[M/X] \models (X(z) \wedge \forall x (X(x) \rightarrow X(u(x))))$. Then $s[M/X](z) \in M$ and whenever $x \in M$, also $(s[M/X](u))(x) \in M$. In other words, since $s[M/X] \sim_X s$, $m_0 \in M$ and if $x \in M$ then $f(x) \in M$, so $m_0 \in M$, $m_1 = f(m_0) \in M$, $m_2 = f(f(m_0)) \in M$, etc. Thus, $M = |\mathfrak{M}|$, and so $\mathfrak{M}, s[M/X] \models \forall x X(x)$. Since $M \subseteq |\mathfrak{M}|$ was arbitrary, we are done: $\mathfrak{M} \models \text{Count}$.

Now assume that $\mathfrak{M} \models \text{Count}$, O.s.,

$$\mathfrak{M}, s \models \forall X ((X(z) \wedge \forall x (X(x) \rightarrow X(u(x)))) \rightarrow \forall x X(x))$$

for some s . Let $m = s(z)$ and $f = s(u)$ and consider $M = \{m, f(m), f(f(m)), \dots\}$. M so defined is clearly **enumerável**. Then

$$\mathfrak{M}, s[M/X] \models (X(z) \wedge \forall x (X(x) \rightarrow X(u(x)))) \rightarrow \forall x X(x)$$

by assumption. Also, $\mathfrak{M}, s[M/X] \models X(z)$ since $M \ni m = s[M/X](z)$, and also $\mathfrak{M}, s[M/X] \models \forall x (X(x) \rightarrow X(u(x)))$ since whenever $x \in M$ also $f(x) \in M$. So, since both antecedent and conditional are satisfied, the consequent must also be: $\mathfrak{M}, s[M/X] \models \forall x X(x)$. But that means that $M = |\mathfrak{M}|$, and so $|\mathfrak{M}|$ is **enumerável** since M is, by definition. $\square$

Problema syn.1. The **sentença** $\text{Inf} \wedge \text{Count}$ is true in all and only **infinito enumerável** domains. Adjust the definition of **Count** so that it becomes a different **sentença** that directly expresses that the domain is **infinito enumerável**, and prove that it does.

Créditos das Imagens

Referências Bibliográficas